arXiv:1303.0451v1 [math.AG] 3 Mar 2013

THE SIGMA FUNCTION FOR WEIERSTRASS SEMIGROUPS $\langle 3, 7, 8\rangle$ AND $\langle 6, 13, 14, 15, 16\rangle$

JIRYO KOMEDA, SHIGEKI MATSUTANI, EMMA PREVIATO

Abstract. Compact Riemann surfaces and their abelian functions are instrumental to solve integrable equations; more recently the representation theory of the Monster and related modular form have pointed to the relevance of τ-functions, which are in turn connected with a specific type of abelian function, the (Kleinian) σ-function. This paper proposes a construction of σ-functions based on the nature of the Weierstrass semigroup at one point of the Riemann surface as a generalization of the construction of plane affine models of the Riemann surface. Because our definition is algebraic, we are able to consider the properties of the σ-functions including their Jacobi inversion formulae, and to give an observation of their properties to those of a Norton basis for replicable functions, in turn relevant to the Monstrous Moonshine.

1. Introduction

We work over the complex field $k = \mathbb{C}$, with compact Riemann surfaces, and simply refer to them as "curves", although in some instances we allow for singular points. We propose a generalization of the σ-function for (n, s) curves, which was developed recently, based on Klein's and Baker's theories [B1]; we parallel constructions given in Mumford's Tata lectures on theta II [Mum], namely algebraic expressions for abelian functions, based on nineteenth-century work; by the use of specific abelian variables and the introduction of a generalized Weierstrass $\wp$-function, defined for hyperelliptic curves [Mum, IIIa. Section 10], Mumford expresses the transcendental function theory on the Jacobian in terms of meromorphic functions on the curve; applications are given to integrable dynamical systems.

1.1. The σ-function.

We say that a pointed algebraic curve is of W-type $(n_1, ..., n_s)$ $(n_1 < n_2 < ... < n_s)$ if its Weierstrass semigroup at the point has minimal set of generators $\{n_1, ..., n_s\}$. The Kleinian σ-function (cf. [B1, B2, K, W]), originally devised for hyperelliptic curves, which have W-type $(2, 2g + 1)$, is closely related to Mumford's $\wp$. Baker showed in essence that σ satisfies the KdV hierarchy and KP equation (as they became known in the 1970s). He introduced a bilinear equation, which was newly proposed in the 1960s as Hirota's bilinear operator [B3]. Baker's work was recently revisited and expanded by several authors, among whom co-authors Buchstaber, Enolskii, and Leykin [BEL1, BEL2, BLE1, BLE2]; Grant [G]; Matsutani [Ma]; Ônishi [O]. In subsequent work [EEL], Enolskii, Eilbeck, and Leykin gave a construction of the Kleinian σ-function of [BLE1] generalized to curves of W-type (n, s), where n and s are any two relatively prime positive integers; we call them (n, s) for short, as in the literature (these are the same as the C_{rs} curves of [MP08, MP12]); They did so by defining the fundamental differential of the second kind over every (n, s) curve. We call their construction "EEL construction". Using the EEL construction, we have investigated properties (such as addition formulae and the order of vanishing) of the σ-function of (n, s) curves in [EEMOP1, EEMOP2, MP08]. Nakayashiki [No] investigated properties of the function also by the use of differentials on symmetric products of the curve. Further Eilbeck1, Enolski and Gibbons [EEG] and Nakayashiki [Na2] showed the connections between σ-function and so-called τ-function [Mul].

In our program, we extend the study to affine curves in higher-dimensional space. We generalize the σ-function to a 3-space curve of W-type $(3, 7, 8)$ and to a curve of W-type $(6, 13, 14, 15, 16)$ which covers it as one of the present authors showed for a curve of W-type $(3, 4, 5)$ in [MK]. As in the (n, s) case, we omit "W-type" for brevity. The motivation of this generalization comes from the the following Monstrous Moonshine problem and Weierstrass semigroup problem. Following the EEL construction, we define the fundamental differential of the second kind, which gives a generalized Legendre relation; this in turn gives us the σ-function and we obtain the Jacobi inversion formulae following previous work [MP08]. We note that our Kleinian σ-function is again a generalization of Weierstrass' elliptic σ.

Korotkin with Shramchenko defined a sigma function for any compact Riemann surface [KS], which so far has not been explicitly related with meromorphic functions on the curve; on the other hand, Ayano investigated sigma functions for space curves of a special type [A], called telescopic curves, but those do not include a (4,7,8) or a (6,13,14,15,16) curve.

1.2. **Monstrous Moonshine.** Our motivation for focusing on the curves of genus 4, $(3, 7, 8)$ and genus 12, $(6, 13, 14, 15, 16)$, is due to a suggestion by John McKay. The curve $(6, 13, 14, 15, 16)$ is associated with a Jacobi variety whose real dimension is 24 and has the Weierstrass gap sequence $\{1, 2, 3, 4, 5, 7, 8, 9, 10, 11, 17, 23\}$, similar to the Norton numbers $\{1, 2, 3, 4, 5, 7, 8, 9, 11, 17, 19, 23\}$. The Norton numbers, as stated in [FMN, McK, MS, No], e.g., are the coefficients of replicable functions in a Norton basis; in turn, replicable functions are related to the Monstrous Moonshine. In [HBJ, VI.3], the **Prize Question** reads, for $\hat{A}$ the (twisted) $\hat{A}$-genus: *Does there exist a 24-dimensional, compact, orientable, differentiable manifold X with $p_1(X) = 0$, $w_2(X) = 0$, $\hat{A}(X) = 1$ and $\hat{A}(X, T_{\mathbb{C}}) = 0$?* For such a manifold, suitable twisted $\hat{A}$ genera would equal dimensions of irreducible representations of the Monster; and if the Monster acted on X by diffeomorphisms, *one would possess a key to construct a great many representations of the Monster.* The related Witten genus is studied in terms of a product of the Weierstrass σ function, there [HBJ, VI.3].

Of course, although a Jacobian is compact, orientable, differentiable and admits a spin structure [Hi] (equivalently, its w_2 is zero), p_1 is non-zero, being the genus of the curve. Since there is no further relation except 24-dimension, we will not be concerned with these problems in this article, but we give some remarks in Section 7 on relevant geometric properties from a viewpoint of the construction of the generalized σ function associated with the curve $(6, 13, 14, 15, 16)$.

With the above motivation, we sought a curve $(6, 13, 14, 15, 16)$. However, as recalled in Section 2, it is a non-trivial problem whether there exists a pointed curve with given numerical semigroup at the point. This article is based upon J. Komeda's proof that there exists a non-singular curve with Weierstrass gap sequence $\{1, 2, 3, 4, 5, 7, 8, 9, 10, 11, 17, 23\}$, at a given point (see Proposition 2.1); its Weierstrass semigroup is indeed H_{12}, the subsemigroup of the non-negative integers $\mathbb{N}_0$ generated by $M_{12} := \{6, 13, 14, 15, 16\}$. The genus-12 curve constructed by Komeda covers a genus-4 curve of $(3, 7, 8)$; we begin with the construction and analysis of its σ-function; relationships between the algebraic functions of the curves appear to be related to the Monstrous Moonshine.

Let H denote a subsemigroup of the additive group of non-negative integers such that only finitely many positive integers, called the gaps of H, are missing from H. Let B_H denote the subring of the polynomial ring $k[t]$ generated by the monomials t^h, $h \in H$. Then Spec B_H is a monomial curve; i.e. an irreducible affine curve with $\mathbb{G}_m$-action, where $\mathbb{G}_m$ denotes the multiplicative group of k ($\mathbb{C}$ in our case). There is an induced $\mathbb{G}_m$-action on $T^1(B_H)$ and, decomposing T^1 into eigenspaces, one can write $T^1 = \sum_\nu T^1(\nu)$, where ν ranges from $-\infty$ to $+\infty$. One says that B_H is negatively graded if $T^1(\nu) = 0$ for all positive ν.

In his thesis [Pi], Pinkham showed that H occurs as a Weierstrass semigroup if and only if the corresponding monomial curve, Spec $k[t^h \colon h \in H]$, can be smoothed negatively. Pinkham used this to show that if H is a complete intersection (in particular, if H is generated by two elements), then H occurs as a Weierstrass semigroup. This also holds for almost complete intersection semigroups generated by three elements and was generalized to almost complete intersection semigroups generated by four elements. Rim and Vitulli [RV] give a complete characterization of such negatively graded semigroup rings. Moreover, they extend Pinkham's result by showing that every negatively graded semigroup ring can be smoothed. By the work of Pinkham, this implies that if H is any negatively graded semi-group, then there exists a smooth projective curve X and a point $x \in X$ such that the gaps of H are the Weierstrass gaps at x.

In Section 2 we give a short review of the Weierstrass Semigroups. In Sections 3 and 4, we proceed as follows: we define monomial curves with given Weierstrass semigroups, with motivation from the Norton numbers; when the semigroup falls outside the verifiably smoothable case, we give Komeda's proof that one smooth curve with such Weierstrass semigroup exists; Pinkham's calculation of the expected dimension yields a positive number, therefore we can conclude that the monomial curve is smoothable. In Section 5, we use the local coordinates given by the monomial presentation of the curve, to manufacture a local section of certain meromorphic differentials, and we construct an abelian function on the curve, the σ-function, by integrating those differentials. In [MK], the original idea was implemented for the semigroup $\langle 3, 4, 5\rangle$, including the σ-function construction and natural extensions of the ones previously developed for (n, s) curves. The σ-function provides a stratification of the Jacobian in Section 6; in Section 7, we make some observations on the possible links with the representation theory of the Monster, starting with the above-mentioned numerical observation connecting the Norton numbers and the differentials we constructed (Appendix A).

We would like to express our thanks to John McKay for posing to us the problem of the Moonshine: his work on replicable functions led him to observations and conjectures (cf. [MS]) which we propose to relate to our work on the σ function; John hosted us at Concordia in 2004 and shared his vision most generously. Further the second author (S.M) has been studied the geometrical interpretations of the replicable function by the guide of John McKay. On the interpretation, Yuji Kodama gave his attentions to the Norton number. John McKay taught him the relations among the Coble's work, sextic curves and E^8 action. The second author is also grateful to Kenichi Tamano, Victor Enolskii, Dmitry Leykin, Yoshihiro Ônishi, Takao Kato and Akira Ohbuchi for valuable discussions and comments.

2. WEIERSTRASS SEMIGROUPS

For reference, we give a brief overview of the recent study of the "numerical semigroups", namely those (additive) sub-semigroups of the non-negative integers $\mathbb{N}_0$ whose complement is finite. We then give a new result due to J. Komeda concerned with the analogy with the Norton basis; this was already presented in [MK].

For a numerical semigroup H generated by a set M, the number of elements of $L(H) := \mathbb{N}_0 \setminus H$, the "gap sequence" of H, is called "genus" if it is finite, and denoted by $g(H)$. We focus on the numerical semigroup H_4 generated by $M_4 := \{3, 7, 8\}$, i.e., $H_4 = \langle 3, 7, 8\rangle$ as well as the problem of the numerical semigroup H_{12} generated by M_{12}, i,e,, $H_{12} = \langle M_{12}\rangle$for the reasons explained in subsection 1.2.

The genus of H_4, H_{12} (respectively) is $g = 4, 12$, and

$$L(H_4) = \{1, 2, 4, 5\}, \quad L(H_{12}) = \{1, 2, 3, 4, 5, 7, 8, 9, 10, 11, 17, 23\}.$$

For a gap sequence $L := \{\ell_0 < \ell_1 < \cdots < \ell_{g-1}\}$ of genus g, let $M(L)$ be the minimal set of generators for the semigroup $H(L)$ (it is easy to show that a minimal set of generators must be unique) and

$$\alpha(L) := \{\alpha_0(L), \alpha_1(L), \dots, \alpha_{g-1}(L)\} \tag{2.1}$$

where $\alpha_i := \ell_i - i - 1$. When an α_i is repeated $j > 1$ times we write α_i^j in $\alpha(L)$. We let $a_{min}(L)$ be the smallest positive integer of $M(L)$. We call a semigroup H an $a_{min}(L)$-semigroup, so for example H_4 is a 3-semigroup and H_{12} is a 6-semigroup and

$$\alpha(L(M_4)) = \{0^2, 1^2\} \quad \text{and} \quad \alpha(L(M_{12})) = \{0^5, 1^5, 6, 11\}.$$

The 6-semigroups with 3 generators, such as H_4, were studied by Komeda in [K04].

For a curve X of genus g and a point $P \in X$, the semigroup

$$H(X,P) := \{n \in \mathbb{N}_0 \mid \text{there exists } f \in k(X) \text{ such that } (f)_\infty = nP\ \},$$

is called the Weierstrass semigroup of the point P. If $L(H(X,P)) := \mathbb{N}_0 \setminus H(X,P)$ differs from the set $\{1, 2, \cdots, g\}$, we say that P is a Weierstrass point of X.

A numerical semigroup H is said to be Weierstrass if there exists a pointed curve (X,P) such that $H = H(X,P)$. Hurwitz posed the problem whether any numerical semigroup H is Weierstrass. The question was revived in the 1980s, viewed as the question of deformations of a reduced complex curve singularity (X_0, ∞). In [BG1, BG2], a counterexample was given, for the semigroup H_B generated by 13, 14, 15, 16, 17, 18, 20, 22 and 23, whose genus is 16: this can be seen as follows. Assume that $H_B = H(X,P)$ for a pointed curve (X,P), then X would have a regular differential vanishing to order $\ell - 1$ at P for any $\ell \in L(H_B)$. By letting $L_2(H)$ be the set of all sums of two elements of $L(H)$ for a semigroup H, X would have a regular quadratic differential vanishing to order $m-2$ at P for any $m \in L_2(H_B)$, which contradicts the fact that $\#L_2(H_B) = 46 > 45 = (3 \times 16) - 3 = h^0(X, \mathcal{K}_X^{\otimes 2})$. Note that we write $\mathcal{K}_X^{\otimes 2}$ for the sheaf of regular quadratic differentials on X and we use standard notation, e.g. h^i for the dimension of the cohomology group H^i of a sheaf over the curve.

We just refer to the work of Komeda for further information; largely thanks to Komeda and co-authors [K83, K99, K04, KO04, KO08, KO08a] and the reference therein, we sketch a list of sufficient properties for H to be Weierstrass, which also includes results by D. Eisenbud with J. Harris, C. Maclachlan and I. Morrison with H. Pinkham,

(1) semigroup whose cardinality of the generators is less than 4,
(2) semigroup whose genus is less than 9,
(3) semigroup whose all primitive sequence, twice the smallest positive integer in $H(L) >$ the largest integer in L, of genus 9,
(4) semigroup whose $a_{\min}(L) = 2, 3, 4, 5$,
(5) semigroup whose $a(L) = (0^{g-2}, m, n)$ for genus g and $wt(L) = g$,
(6) semigroup whose $a(L) = (0^{g-r}, m^r)$ for genus g, and
(7) other many special cases of n-semigroups generated by 4 elements.

As it is partially mentioned in [MK], we have the following crucial proposition due to Komeda:

Proposition 2.1. *The numerical semigroup $\langle 6, 13, 14, 15, 16\rangle$ is Weierstrass.*

Proof. We consider a pointed curve (X,P) with $H(X,P) = \langle 3, 7, 8\rangle$ and $\mathcal{K}_X$ a canonical divisor on X. Then,

$$2 = h^0(4P) = 4 + 1 - 4 + h^0(\mathcal{K}_X - 4P) = 1 + h^0(\mathcal{K}_X - 4P);$$

this implies that $\mathcal{K}_X - 4P \sim P_1 + P_2$ for some points P_1 and $P_2 \in X$. Moreover, since

$$2 = h^0(5P) = 5 + 1 - 4 + h^0(\mathcal{K}_X - 5P) = 2 + h^0(\mathcal{K}_X - 5P),$$

then $h^0(\mathcal{K}_X - 5P) = 0$, and $P_i \neq P$ for $i = 1, 2$. This implies that $\mathcal{K}_X \sim 4P + P_1 + P_2$ with $P_i \neq P$ for $i = 1, 2$. By setting $D := 7P - P_1 - P_2$, we have $\deg(2D - P) = 9 = 2 \times 4 + 1$, and this implies that the complete linear system $|2D - P|$ is very ample, hence base-point free. Therefore, $2D \sim P + Q_1 + \ldots + Q_9$ (= a reduced divisor). Let $\mathcal{L}$ be the invertible sheaf $\mathcal{O}_X(-D)$ on X and ϕ an isomorphism $\mathcal{L}^{\otimes 2} \approx \mathcal{O}_X(-P - Q_1 - \cdots - Q_9) \subset \mathcal{O}_X$. The vector bundle $\mathcal{O}_X \oplus \mathcal{L}$ has an $\mathcal{O}_X$-algebra structure through ϕ. The canonical $2:1$ morphism $\pi : \tilde{X} = \mathrm{Spec}\ (\mathcal{O}_X \oplus \mathcal{L}) \to X$ has branch locus $\{P, Q_1, \ldots, Q_9\}$. For $\tilde{P}$ the ramification point of π over P, we see that $H(\tilde{P}) = \langle 6, 13, 14, 15, 16 \rangle$ using the following formula:

$$h^0(2n\tilde{P}) = h^0(nP) + h^0(nP - D)$$

for any non-negative integer n. We have $h^0(12\tilde{P}) = h^0(6P) + h^0(6P - 7P + P_1 + P_2) = 3 + h^0(P_1 + P_2 - P) = 3$ because of $P_i \neq P$ for $i = 1, 2$. Moreover, we have $h^0(14\tilde{P}) = h^0(7P) + h^0(P_1 + P_2) = 4 + 1 = 5$, which implies that $13, 14 \in H(\tilde{P})$. The equalities

$$h^0(16\tilde{P}) = h^0(8P) + h^0(P + P_1 + P_2) = 5 + 3 + 1 - 4 + h^0(\mathcal{K}_X - P - P_1 - P_2) = 5 + h^0(3P) = 7$$

mean that $15, 16 \in H(\tilde{P})$. We have

$$h^0(18\tilde{P}) = h^0(9P) + h^0(2P + P_1 + P_2) = 6 + 4 + 1 - 4 + h^0(\mathcal{K}_X - 2P - P_1 - P_2) = 7 + h^0(2P) = 8,$$

which implies that $17 \notin H(\tilde{P})$ and $18 \in H(\tilde{P})$. Moreover, we get

$$h^0(20\tilde{P}) = h^0(10P) + h^0(3P + P_1 + P_2) = 7 + 5 + 1 - 4 + h^0(\mathcal{K}_X - 3P - P_1 - P_2) = 9 + h^0(P) = 10.$$

Hence, we have $19, 20 \in H(\tilde{P})$. We obtain

$$h^0(22\tilde{P}) = h^0(11P) + h^0(4P + P_1 + P_2) = 8 + 6 + 1 - 4 + h^0(\mathcal{K}_X - 4P - P_1 - P_2) = 11 + 1 = 12.$$

Hence, we have $21, 22 \in H(\tilde{P})$. Finally, we get

$$h^0(24\tilde{P}) = h^0(12P) + h^0(5P + P_1 + P_2) = 9 + 7 + 1 - 4 = 13.$$

Hence, we have $23 \notin H(\tilde{P})$ and $24 \in H(\tilde{P})$. We conclude that $H(\tilde{P}) = \langle 6, 13, 14, 15, 16 \rangle$. □

3. SEMIGROUP $H = \langle 3, 7, 8 \rangle$

In our construction of σ-functions, the tool is a description of the affine ring of the curve minus ∞, the Weierstrass point in question. We begin with semigroup $H = \langle 3, 7, 8 \rangle$ because to treat our motivating example H_{12}, the existence proof in Proposition 2.1, we use an explicit double cover of a curve with semigroup H_4.

3.1. **The curve as a monomial curve.** A monomial curve is an irreducible affine curve with $\mathbb{G}_m$-action, where $\mathbb{G}_m$ is the multiplicative group of the complex numbers. To construct our curve, we consider the semigroup ring of H_4. We follow Pinkham's strategy [Pi] both in the H_4 and H_{12} cases, namely, we start with an irreducible curve singularity with $\mathbb{G}_m$ action, described parametrically by its semigroup ring. We deform the matrix of the relations. This deformation space $\bar{U}$ "classifies the set of pairs consisting of a smooth and proper algebraic curve X together with a point $P \in X$ with given semigroup $H(X, P)$. If $\bar{U}$ is non void, then we have constructed directly a compactification of a "moduli space" for curves with points of semi group H" [Pi, Introduction]. We have a result that ensures that $\bar{U}$ is nonempty in both cases. In the case of H_4, Pinkham again [Pi, Section 14] proves nonemptiness for semigroups of two generators and conjectures it for three, proved later in [RV].

Proposition 3.1. *For the k-algebra homomorphism,*

$$\varphi_4 : k[Z] := k[Z_3, Z_7, Z_8] \to k[t^a]_{a\in M_4}, \quad Z_a \mapsto t^a,$$

the kernel of φ_4 is generated by $f^{(Z)}_b = 0$ $(b = 14, 15, 16)$ where

$$f^{(Z)}_{14} = Z_7^2 - Z_3^2 Z_8, \quad f^{(Z)}_{15} = Z_7 Z_8 - Z_3^5, \quad f^{(Z)}_{16} = Z_8^2 - Z_3^3 Z_7. \tag{3.1}$$

Proof. This follows from a result of Herzog's [Her], who showed that for any W-semigroup, the number of generators of the kernel is two or three. □

Here we note that these relations are given by the 2×2 minors of

$$\begin{vmatrix} Z_3^2 & Z_7 & Z_8 \\ Z_7 & Z_8 & Z_3^3 \end{vmatrix}. \tag{3.2}$$

The monomial ring $B_H := k[t^a]_{a=M_4}$ is given by

$$B_H \simeq k[Z_3, Z_7, Z_8]/\mathrm{ker}\varphi.$$

We say that Z_a has weight a.

To indicate the action of $\mathbb{G}_m$, defined as $Z_a \mapsto g^a Z_a$, and the induced action on the monomial ring B_{H_4}, we use the notation $\mathbb{G}_m^{(4)}$: it will be convenient since we are going to relate several curves and different actions.

To construct a smooth curve X_4 with $H(X_4, \infty) = H_4$, we consider two plane affine curves X_6 and X_7, with infinity, corresponding to the equations:

$$f_{6,21}(x, y_7) := y_7^3 - k_7(x), \quad k_7(x) := k_3(x)k_2(x)^2,$$

and

$$f_{7,24}(x, y_8) := y_8^3 - k_8(x), \quad k_8(x) := k_3(x)^2 k_2(x),$$

where for distinct $b_a \in \mathbb{C}$ $(a = 1, 2, \cdots, 5)$,

$$k_3(x) := (x - b_1)(x - b_2)(x - b_3) = x^3 + \lambda^{(3)}_1 x^2 + \lambda^{(3)}_2 x + \lambda^{(3)}_3,$$

$$k_2(x) := (x - b_4)(x - b_5) = x^2 + \lambda^{(2)}_1 x + \lambda^{(2)}_2.$$

Both curves have the singular points in their affine parts; the singular points of the affine part of X_6 are $(b_a, 0)$ $a = 4, 5$ whereas those of X_7 are $(b_a, 0)$ $a = 1, 2, 3$. We consider the rings $R_6 := \mathbb{C}[x, y_7]/(f_{6,21}(x, y_7))$ and $R_7 := \mathbb{C}[x, y_8]/(f_{7,24}(x, y_8))$ associated with X_6 and X_7 respectively. The genera of the semigroups of X_6 and X_7 at their ∞ are $6 = g(\langle 3, 7\rangle)$ and $7 = g(\langle 3, 8\rangle)$ respectively.

Due to the normalization theorem [Gri, p.5, p.68] (Theorem B.1), it is known that every algebraic curve (due to its dimensionality) admits a normalization and the normalization is unique up to isomorphism. Although the functions y_7 and y_8 we introduced are defined on different curves, we consider that the following identities are consistent because the zeroes of $f_{6,21}(x, y_7)$ and $f_{7,24}(x, y_8)$ satisfy the relations,

$$y_7 y_8 = k_2(x)k_3(x), \quad y_8 = \frac{y_7^2}{(x - b_4)(x - b_5)}, \quad y_7 = \frac{y_8^2}{(x - b_1)(x - b_2)(x - b_3)}, \tag{3.3}$$

where consistency is achieved by requiring that under the natural action of a primitive third root of unity ζ_3 on y_7 and y_8 in R_6 and R_7 as respectively: on the first relation, we make the choice $a = 0$ out of the possible $0 \le a \le 2$ of $y_7 y_8 \mapsto \zeta_3^a k_2(x)k_3(x)$.

As a normalisation of these singular curves, we consider the commutative ring,

$$R_4 := \mathbb{C}[x, y_7, y_8]/(f_{14}, f_{15}, f_{16}),$$

and the curve Spec R_4. Here we define $f_{14}, f_{15}, f_{16} \in \mathbb{C}[x, y_7, y_8]$ by

$$f_{14} = y_7^2 - y_8 k_2(x), \quad f_{15} = y_7 y_8 - k_2(x)k_3(x), \quad f_{16} = y_8^2 - y_7 k_3(x),$$

which are also viewed as the 2×2 minors of

$$\begin{vmatrix} k_2(x) & y_7 & y_8 \\ y_7 & y_8 & k_3(x) \end{vmatrix}, \tag{3.4}$$

as a deformation of (3.2). Note that when x is infinite, so are y_7 and y_8. $\mathbb{G}_m^{(4)}$ acts on R_4 by sending x and y_a to $g^{-3}x$, $g^{-a}y_a$ for $g \in \mathbb{G}_m^{(4)}$ and $a = 7, 8$ at $\infty \in X_4$. Thus -3 and $-a$ are the weights of x and y_a at ∞.

Due to Nagata's Jacobi criterion [Mat, Theorem 30.10], Spec R_4 is non-singular except at ∞:

Proposition 3.2. *For every (x, y_7, y_8) where f_{14}, f_{15} and f_{16} simultaneously vanish,*

$$\operatorname{rank} \mathcal{U}_4 = 2,$$

where

$$\mathcal{U}_4 := \begin{pmatrix} \frac{\partial f_{14}}{\partial x} & \frac{\partial f_{14}}{\partial y_7} & \frac{\partial f_{14}}{\partial y_8} \\ \frac{\partial f_{15}}{\partial x} & \frac{\partial f_{15}}{\partial y_7} & \frac{\partial f_{15}}{\partial y_8} \\ \frac{\partial f_{16}}{\partial x} & \frac{\partial f_{16}}{\partial y_7} & \frac{\partial f_{16}}{\partial y_8} \end{pmatrix}.$$

Proof. The matrix $\mathcal{U}_4$ is equal to

$$\mathcal{U}_4 = \begin{pmatrix} y_8 k_2'(x) & 2y_7 & -k_2(x) \\ (k_2(x)k_3(x))' & y_8 & y_7 \\ y_7 k_3'(x) & -k_3(x) & 2y_8 \end{pmatrix}.$$

When y_7 vanishes, and in consequence y_8 vanishes, x must be one of $\{b_a\}_{a=1,\cdots,5}$. For $x = b_1$,

$$\mathcal{U}_4 = \begin{pmatrix} 0 & 0 & -k_2(b_1) \\ k_2(b_1)(b_1 - b_2)(b_1 - b_3) & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}.$$

Here $k_2(b_1)$ and $(b_1 - b_2)(b_1 - b_3)$ don't vanish so the rank is 2. Similarly for $x = b_a$ $a = 2, \cdots, 5$. On the other hand, when y_7 and y_8 are nonzero,

$$\begin{pmatrix} 1 & & \\ 1 & -1 & 1 \\ & & 1 \end{pmatrix} \begin{pmatrix} y_8^2 & & \\ & y_7 y_8 & \\ & & y_7^2 \end{pmatrix} \mathcal{U}_4 = \begin{pmatrix} 1 & & \\ 1 & -1 & 1 \\ & & 1 \end{pmatrix} \begin{pmatrix} y_8^3 k_2'(x) & 2y_7 y_8^2 & -y_8^2 k_2(x) \\ y_7 y_8 (k_2(x)k_3(x))' & y_7 y_8^2 & y_7^2 y_8 \\ y_7^3 k_3'(x) & -y_7^2 k_3(x) & 2y_7^2 y_8 \end{pmatrix}$$

shows that the rank is 2. ☐

Lastly, the smooth curve X_4 associated to the quotient field of the ring R_4 is obtained by completing the affine piece Spec R_4 by one smooth point at infinity, which we still denote by ∞ when there is no ambiguity; this can be seen by introducing a local parameter at ∞ as in [Mul, Section 2] (see Appendix B). There is a cyclic action on X_4 by

$$\hat{\zeta}_3(x, y_7, y_8) = (x, \zeta_3 y_7, \zeta_3^2 y_8).$$

Proposition 3.3. *There is a natural homomorphism $R_a \to R_4$ for each $(a = 6, 7)$.*

Proof. Letting $\varphi : \mathbb{C}[x, y_7] \to \mathbb{C}[x, y_7, y_8]$ be a ring homomorphism, we have the relation $\varphi^{-1}(f_{14}, f_{15}, f_{16}) = (f_{6,21})$. Similarly we have the result for $a = 8$. ☐

In conclusion, X_4 is the normalization of X_6 and X_7.

3.2. **The Weierstrass gaps and holomorphic one forms.** The Weierstrass gap sequences at ∞ are given by the following table. Indeed, we have $H(X_4, \infty) = H_4$. In view of the construction of the curve, functions that have the order of poles in the complement of the gaps can be given by monomials in R_4. We denote by $\phi_i^{(g)}$ a monomial in R_g for $g = 6, 7$ and we add the notational device H^0 in the subscript to signify functions, as in the group $H^0(X_4 \backslash \infty, \mathcal{O}_{X_4})$: $\phi_{H^0 j}^{(4)}$ for R_4, *e.g.*, $\phi_{H^0 0}^{(4)} = 1$, $\phi_{H^0 1}^{(4)} = x$, $\phi_{H^0 2}^{(4)} = x^2$, $\phi_{H^0 3}^{(4)} = y_7$, $\phi_{H^0 4}^{(4)} = y_8$, $\phi_{H^0 5}^{(4)} = x^3$, e.g..

Table 1

	0	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16
X_6	1	-	-	x	-	-	x^2	y_7	-	x^3	xy_7	-	x^4	x^2y_7	y_7^2	x^5	x^3y_7
X_7	1	-	-	x	-	-	x^2	-	y_8	x^3	-	xy_8	x^4	-	x^2y_8	x^5	y_8^2
X_4	1	-	-	x	-	-	x^2	y_7	y_8	x^3	xy_7	xy_8	x^4	x^2y_7	x^2y_8	y_7y_8	x^3y_7

We define the weight $N^{(g)}(n)$ and $N^{(4)}_{H^0}(n)$ by

$$N^{(4)}_{H^0}(n) = -\mathrm{wt}(\phi^{(4)}_{H^0 n}), \quad N^{(g)}(n) = -\mathrm{wt}(\phi^{(g)}_n),$$

where wt() is the order of pole at ∞. These are consistent with the weights corresponding to the action $\mathbb{G}_m^{(4)}$, whereas R_g $(g = 6, 7)$ provides the Weierstrass gap sequences associated with the semigroups $\langle 3, 7\rangle$ and $\langle 3, 8\rangle$.

The differentials of the first kind of the singular curves of X_6 and X_7 are given by,

$$\nu_i^{I;6} := \frac{\phi^{(7)}_{i-1}dx}{3y_7^2}, \quad (i = 1, 2, 3, 4, 5, 6), \quad \nu_i^{I;7} := \frac{\phi^{(8)}_{i-1}dx}{3y_8^2}, \quad (i = 1, 2, 3, 4, 5, 6, 7).$$

Proposition 3.4. *The holomorphic one form over X_4 can be expressed by*

$$\nu_i^{I;4} := \frac{\phi^{(4)}_{H^1 i-1}dx}{3y_7y_8}, \quad (i = 1, 2, 3, 4),$$

where the monomial $\phi^{(4)}_{H^1 i} \in R_4$ is defined by

$$\phi^{(4)}_{H^1 0} := y_7, \quad \phi^{(4)}_{H^1 1} := y_8, \quad \phi^{(4)}_{H^1 2} := xy_7, \quad \phi^{(4)}_{H^1 3} := xy_8$$

and we add the notational device H^1 in the subscript to indicate differentials, as in the group $H^1(X_4\backslash\infty, \mathcal{O}_{X_4})$:

Further we define $\phi^{(4)}_{H^1 4} := x^2y_7$, $\phi^{(4)}_{H^1 5} := x^2y_8$, and for $j > 5$, and let $\phi^{(4)}_{H^1 j}$ defined as $\phi^{(4)}_{H^1 j} := \phi^{(4)}_{H^0 j+5}$. We also define the weight $N^{(4)}_{H^1}(n)$ by

$$N^{(4)}_{H^1}(n) = -\mathrm{wt}(\phi^{(4)}_{H^1 n}),$$

the order of pole of $\phi^{(4)}_{H^1 n}$ at ∞, which is also consistent with the weight coming from $\mathbb{G}_m^{(4)}$.

Remark 3.5. We remark the canonical divisor $\mathcal{K}_{X_4}$ of X_4. Since by letting a point in X_4 by (x, y_7, y_8), the divisors of the bases are

$$\begin{aligned}
(\nu_1^{I;4}) &= (b_4, 0, 0) + (b_5, 0, 0) + 4\infty, \\
(\nu_2^{I;4}) &= (b_1, 0, 0) + (b_2, 0, 0) + (b_3, 0, 0) + 3\infty, \\
(\nu_3^{I;4}) &= \sum_{a=0}^{2}(0, \zeta_3^a\sqrt{3}k_7(0), \zeta_3^{2a}\sqrt{3}k_8(0)) + (b_4, 0, 0) + (b_5, 0, 0), \\
(\nu_4^{I;4}) &= \sum_{a=0}^{2}(0, \zeta_3^a\sqrt{3}k_7(0), \zeta_3^{2a}\sqrt{3}k_8(0)) + (b_1, 0, 0) + (b_2, 0, 0) + (b_3, 0, 0),
\end{aligned}$$

we emphasise that the canonical divisor $\mathcal{K}_{X_4}$ must include points in $X_4 \setminus \infty$. It shows a contrast to the fact that for every (r, s) curve of genus $g = (r-1)(s-1)/2$, there is a canonical divisor given by $(2g-2)\infty$ as in [MP09]. This is a reason why we consider $\phi_{H^1}^{(4)}$'s besides $\phi_{H^0}^{(4)}$'s.

Then the following proposition is self-evident:

Proposition 3.6. *For $n > 0$, $\sum_{i=0}^n a_i \frac{\phi_{H^1\,i}^{(4)} dx}{y_7 y_8}$ belongs to $H^0(X_4 \setminus \infty, \Omega_{X_4})$, in particular to $H^0(X_4, \Omega_{X_4})$ for $n \le 4$, where Ω_{X_4} is the sheaf of the holomorphic one-form.*

Lemma 3.7. *For a non-negative integer $n < 5$, if the one form*

$$\sum_{i=0}^{n} a_i \frac{x^i dx}{y_7 y_8} \equiv \sum_{i=0}^{n} a_i \frac{x^i dx}{k_2(x) k_3(x)}$$

is holomorphic over X_4, then each a_i vanishes.

Proof. For $n < 5$, every term in $\sum_{i=0}^n a_i \frac{x^i dx}{y_7 y_8}$ has singularities at points in $X \setminus \infty$. □

By letting

$$\Lambda_i^{(g)} = N^{(g)}(g) - N^{(g)}(i-1) - g + i - 1, \qquad \Lambda_i^{(4)} = N_{H^1}^{(4)}(4) - N_{H^1}^{(4)}(i-1) - 5 + i,$$

the related Young diagrams $\mathcal{Y}_4 = (\Lambda_1^{(4)}, \Lambda_2^{(4)}, \Lambda_3^{(4)}, \Lambda_4^{(4)})$, $\mathcal{Y}_6 = (\Lambda_1^{(6)}, \Lambda_2^{(6)}, \cdots, \Lambda_6^{(6)})$ and $\mathcal{Y}_7 = (\Lambda_1^{(7)}, \Lambda_2^{(7)}, \cdots, \Lambda_7^{(7)})$ are given by

 ,  ,  .

Since the Young diagram $\mathcal{Y}_4$ is not symmetric, the semigroup H_4 is also not symmetric, where a numerical semigroup is symmetric if and only if the gap sequence has $2g-1$ [RG]. We should note that $\alpha_{g-i}(L)$ in (2.1) corresponds to $\Lambda_i^{(g)}$ by appropriate interpretations for each X_g $(g = 4, 6, 7)$.

Also self-evident is the following:

Lemma 3.8. *By using the same letters in source and target under the homomorphism of Proposition 3.3 (as well as identifying a coset by a standard representative), the holomorphic one forms* $\nu_i^{I;4}$ *over* X_4 *satisfy:*

$$\nu_i^{I;4} = \tilde{\nu}_{i+3}^{I;6} = \tilde{\nu}_{i+4}^{I;7}, \quad (i = 1, 2, 3, 4),$$

where

$$\tilde{\nu}_i^{I;6} := \nu_i^{I;6}, \quad (i = 4, 6), \quad \tilde{\nu}_3^{I;6} := \nu_3^{I;6} + \lambda_1^{(2)} \nu_2^{I;6} + \lambda_2^{(2)} \nu_1^{I;6}, \quad \tilde{\nu}_5^{I;6} := \nu_5^{I;6} + \lambda_1^{(2)} \nu_3^{I;6} + \lambda_2^{(2)} \nu_2^{I;6},$$

$$\tilde{\nu}_i^{I;7} := \nu_i^{I;7}, \quad (i = 4, 6), \quad \tilde{\nu}_5^{I;7} := \nu_5^{I;7} + \lambda_1^{(3)} \nu_3^{I;7} + \lambda_2^{(3)} \nu_2^{I;7} + \lambda_3^{(3)} \nu_1^{I;7},$$

$$\tilde{\nu}_7^{I;7} := \nu_7^{I;7} + \lambda_1^{(3)} \nu_5^{I;7} + \lambda_2^{(3)} \nu_3^{I;7} + \lambda_3^{(3)} \nu_2^{I;7}.$$

As is standard practice, we choose a basis $\alpha_i^{(4)}, {\beta^{(4)}}_j$ $(1 \leqq i, j \leqq 4)$ of $H_1(X, \mathbb{Z})$ such that their intersection numbers are $\alpha_i^{(4)} \cdot \alpha_j^{(4)} = {\beta^{(4)}}_i \cdot {\beta^{(4)}}_j = 0$ and $\alpha_i^{(4)} \cdot {\beta^{(4)}}_j = \delta_{ij}$, and we denote the period matrices by

$$\left[\, \omega^{(4)\prime} \; \omega^{(4)\prime\prime} \right] = \frac{1}{2} \left[\int_{\alpha_i^{(4)}} \nu_j^{I;4} \quad \int_{{\beta^{(4)}}_i} \nu_j^{I;4} \right]_{i,j=1,2,\cdots,4}. \tag{3.5}$$

Let Λ_4 be a lattice generated by $\omega^{(4)\prime}$ and $\omega^{(4)\prime\prime}$. For a point $P \in X_4$, the Abel map is defined,

$$\hat{u}(P) = \int_\infty^P \nu^{I;4} \in \mathbb{C}^4,$$

and for k points $P_1, P_2, \dots, P_k \in X_4$, the Abel map $\hat{u}(P_1, \dots, P_k)$ is defined by $\sum_{i=1}^k \hat{u}(P_i)$. Then we obtain the Jacobian $\mathcal{J}_4$ by

$$\kappa : \mathbb{C}^4 \to \mathcal{J}_4 = \mathbb{C}^4/\Lambda_4,$$

and the subvariety ${\mathcal{W}^{(4)}}_k$ by

$${\mathcal{W}^{(4)}}_k := \kappa\hat{u}(S^k X_4).$$

Hereafter we use the convention that for $P_a \in X_4$, P_a is expressed by $(x_a, y_{7,a}, y_{8,a})$ or $(x_{P_a}, y_{7,P_a}, y_{8,P_a})$. By letting $u := \hat{u}(P_1, \cdots, P_4)$, we have

$$\begin{pmatrix} \partial/\partial u_1 \\ \partial/\partial u_2 \\ \partial/\partial u_3 \\ \partial/\partial u_4 \end{pmatrix} = \Psi_4^{(4)-1} \begin{pmatrix} 3y_{7,1}y_{8,1}\partial/\partial x_1 \\ 3y_{7,2}y_{8,2}\partial/\partial x_2 \\ 3y_{7,3}y_{8,3}\partial/\partial x_3 \\ 3y_{7,4}y_{8,4}\partial/\partial x_4 \end{pmatrix}, \tag{3.6}$$

where

$$\Psi_4^{(4)} := \begin{pmatrix} \phi^{(4)}_{H^1 0}(P_1) & \phi^{(4)}_{H^1 1}(P_1) & \phi^{(4)}_{H^1 2}(P_1) & \phi^{(4)}_{H^1 3}(P_1) \\ \phi^{(4)}_{H^1 0}(P_2) & \phi^{(4)}_{H^1 1}(P_2) & \phi^{(4)}_{H^1 2}(P_2) & \phi^{(4)}_{H^1 3}(P_2) \\ \phi^{(4)}_{H^1 0}(P_3) & \phi^{(4)}_{H^1 1}(P_3) & \phi^{(4)}_{H^1 2}(P_3) & \phi^{(4)}_{H^1 3}(P_3) \\ \phi^{(4)}_{H^1 0}(P_4) & \phi^{(4)}_{H^1 1}(P_4) & \phi^{(4)}_{H^1 2}(P_4) & \phi^{(4)}_{H^1 3}(P_4) \end{pmatrix}. \tag{3.7}$$

In other words,

$$\sum_{i=1} \epsilon_i \frac{\partial}{\partial u_i} = |\Psi_4^{(4)-1}| \begin{vmatrix} \phi^{(4)}_{H^1 0}(P_1) & \phi^{(4)}_{H^1 1}(P_1) & \phi^{(4)}_{H^1 2}(P_1) & \phi^{(4)}_{H^1 3}(P_1) & 3y_{7,1}y_{8,1}\partial/\partial x_1 \\ \phi^{(4)}_{H^1 0}(P_2) & \phi^{(4)}_{H^1 1}(P_2) & \phi^{(4)}_{H^1 2}(P_2) & \phi^{(4)}_{H^1 3}(P_2) & 3y_{7,2}y_{8,2}\partial/\partial x_2 \\ \phi^{(4)}_{H^1 0}(P_3) & \phi^{(4)}_{H^1 1}(P_3) & \phi^{(4)}_{H^1 2}(P_3) & \phi^{(4)}_{H^1 3}(P_3) & 3y_{7,3}y_{8,3}\partial/\partial x_3 \\ \phi^{(4)}_{H^1 0}(P_4) & \phi^{(4)}_{H^1 1}(P_4) & \phi^{(4)}_{H^1 2}(P_4) & \phi^{(4)}_{H^1 3}(P_4) & 3y_{7,4}y_{8,4}\partial/\partial x_4 \\ \epsilon_1 & \epsilon_2 & \epsilon_3 & \epsilon_4 & \end{vmatrix}.$$

As these relations hold for the image of the Abel map, we also have similar relations for a stratum $\kappa\hat{u}(S^k X_4)$ of $k < 4$ using the submatrices of $\Psi_4^{(4)}$ as in [MP12]. Further we have a natural relation:

$$\sum_{i,j=1}^{4} \phi^{(4)}_{H^1 i-1}(P_1)\phi^{(4)}_{H^1 j-1}(P_2) \frac{\partial^2}{\partial \hat{u}_i(P_1)\partial \hat{u}_j(P_2)} = 9y_{7,1}y_{8,1}y_{7,2}y_{8,2} \frac{\partial^2}{\partial x_1 \partial x_2}. \tag{3.8}$$

3.3. **Differentials of the second and the third kinds.** Following the EEL-construction [EEL, BEL2] for (n,s) curves, we produce an algebraic representation of a differential form which, up to a tensor of holomorphic one-forms, is equal to the fundamental normalized differential of the second kind in [Fay, Corollary 2.6]. We extend the EEL-construction to the space curve X_4.

Definition 3.9. *A two-form $\Omega^{(4)}(P_1, P_2)$ on $X_4 \times X_4$ is called a fundamental differential of the second kind if it is symmetric,*

$$\Omega^{(4)}(P_1, P_2) = \Omega^{(4)}(P_2, P_1), \tag{3.9}$$

it has its only pole (of second order) along the diagonal of $X_4 \times X_4$, and in the vicinity of each point (P_1, P_2) is expanded in power series as

$$\Omega^{(4)}(P_1, P_2) = \Big(\frac{1}{(t_{P_1} - t'_{P_2})^2} + d_{\geq}(1)\Big) dt_{P_1} \otimes dt_{P_2} \quad (\text{as } P_1 \to P_2), \tag{3.10}$$

where t_P is a local coordinate at the point $P \in X_4$.

Proposition 3.10. *Let $\Sigma^{(4)}\big(P_1, P_2\big)$ be the following form,*

$$\Sigma^{(4)}\big(P, Q\big) := \frac{y_{7,P}y_{8,P} + y_{7,P}y_{8,Q} + y_{7,Q}y_{8,P}}{(x_P - x_Q)3y_{7,P}y_{8,P}} dx_P. \tag{3.11}$$

Then $\Sigma^{(4)}(P, Q)$ has the properties

(1) $\Sigma^{(4)}(P,Q)$ as a function of P is singular at $Q = (x_Q, y_{7,Q}, y_{8,Q})$ and ∞, and vanishes at $\hat{\zeta}_3(Q) = (x_Q, \zeta_3 y_{7,Q}, \zeta_3^2 y_{8,Q})$.
(2) $\Sigma^{(4)}(P,Q)$ as a function of Q is singular at P and at ∞.

Proof. Direct computation. $\square$

The following holds for non-singular X_4:

Proposition 3.11. *There exist differentials $\nu_j^{II;4} = \nu_j^{II;4}(x,y)$ $(j = 1, 2, \cdots, 4)$ of the second kind such that they have a simple pole at ∞ and satisfy the relation,*

$$\begin{aligned} &d_Q\Sigma^{(4)}\big(P,Q\big) - d_P\Sigma^{(4)}\big(Q,P\big) \\ &\qquad = \sum_{i=1}^{4}\Big(\nu_i^{I;4}(Q)\otimes\nu_i^{II;4}(P) - \nu_i^{I;4}(P)\otimes\nu_i^{II;4}(Q)\Big), \end{aligned} \tag{3.12}$$

where

$$d_Q\Sigma^{(4)}\big(P,Q\big) := dx_P \otimes dx_Q \frac{\partial}{\partial x_Q}\frac{y_{7,P}y_{8,P} + y_{7,P}y_{8,Q} + y_{7,Q}y_{8,P}}{(x_P - x_Q)3y_{7,P}y_{8,P}} dx_P. \tag{3.13}$$

The set of differentials $\{\nu_1^{II;4},\ \nu_2^{II;4},\ \nu_3^{II;4},\ \nu_4^{II;4}\}$ is determined modulo the linear space spanned by $\langle\nu_j^{I;4}\rangle_{j=1,\dots,4}$ and it has representatives

$$\begin{aligned} \nu_4^{II;4} &= \frac{-x^2dx}{3y_8}, \\ \nu_3^{II;4} &= \frac{-(2x^2 + \lambda_1^{(2)}x)dx}{3y_7}, \\ \nu_2^{II;4} &= \frac{-\Big(4x^3 + (3\lambda_1^{(3)} + 2\lambda_1^{(2)})x^2 + (2\lambda_2^{(3)} + \lambda_1^{(2)}\lambda_1^{(3)})x + \lambda_3^{(3)}\Big)\,dx}{3y_8}, \\ \nu_1^{II;4} &= \frac{-\Big(5x^3 + (3\lambda_1^{(3)} + 4\lambda_1^{(2)})x^2 + (\lambda_2^{(3)} + 2\lambda_1^{(2)}\lambda_1^{(3)} + 3\lambda_2^{(2)})x + \lambda_2^{(2)}\lambda_1^{(3)}\Big)\,dx}{3y_7}. \end{aligned}$$

We will henceforth use this basis $\nu_i^{II;4}$.

Proof.

$$\begin{aligned} &\frac{\partial}{\partial x_Q}\frac{y_{7,P}y_{8,P} + y_{7,P}y_{8,Q} + y_{7,Q}y_{8,P}}{(x_P - x_Q)3y_{7,P}y_{8,P}}dx_P \\ &= \frac{1}{(x_P - x_Q)9y_{7,P}y_{8,P}y_{7,Q}y_{8,Q}}\Big[\frac{3(y_{7,P}y_{8,P} + y_{7,P}y_{8,Q} + y_{7,Q}y_{8,P})y_{7,Q}y_{8,Q})}{(x_P - x_Q)} \\ &\qquad + \Big(y_{7,P}\frac{y_{7,Q}}{y_{8,Q}}(2k_{3,Q}k'_{3,Q}k_{2,Q} + k_{3,Q}^2k'_{2,Q}) - y_{8,P}\frac{y_{8,Q}}{y_{7,Q}}(2k_{3,Q}k_{7,Q}k'_{7,Q} + k'_{3,Q}k_{7,Q}^2))\Big)\Big]. \end{aligned}$$

Here we set $k_{a,P} = k_a(x_P)$ and $k'_{a,P} = dk_a(x_P)/dx_P$. The result follows from the equalities

$$\frac{\partial}{\partial x_Q}\frac{y_{7,P}y_{8,P} + y_{7,P}y_{8,Q} + y_{7,Q}y_{8,P}}{(x_P - x_Q)3y_{7,P}y_{8,P}} - \frac{\partial}{\partial x_P}\frac{y_{7,Q}y_{8,Q} + y_{7,Q}y_{8,P} + y_{7,P}y_{8,Q}}{(x_Q - x_P)3y_{7,Q}y_{8,Q}}$$

is equal to

$$\begin{aligned}
&= \frac{1}{(x_P - x_Q)9y_{7,P}y_{8,P}y_{7,Q}y_{8,Q}}\Big[\frac{3(y_{7,P}y_{8,Q} + y_{7,Q}y_{8,P})(y_{7,Q}y_{8,Q} - y_{7,P}y_{8,P})}{(x_P - x_Q)} \\
&\quad -\Big(y_{7,P}y_{8,Q}(2k'_{3,Q}k_{2,Q} + k_{3,Q}k'_{2,Q} - 2k_{3,P}k'_{2,P} + k'_{3,P}k_{2,P}) \\
&\qquad + y_{7,Q}y_{8,P}(2k'_{3,P}k_{2,P} + k_{3,P}k'_{2,P} - 2k_{3,Q}k'_{2,Q} + k'_{3Q}k_{2,Q})\Big)\Big] \\
&= \frac{1}{(x_P - x_Q)9y_{7,P}y_{8,P}y_{7,Q}y_{8,Q}}\left(B_4(P,Q) - B_4(Q,P)\right).
\end{aligned}$$

where

$$\begin{aligned}
B_4(P,Q) = y_{7,P}y_{8,Q}\Big(&5x_Q^3 + (3\lambda_1^{(3)} + 4\lambda_1^{(2)})x_Q^2 + (\lambda_2^{(3)} + 2\lambda_1^{(2)}\lambda_1^{(3)} + 3\lambda_2^{(2)})x_Q \\
&+ (2x_Q^2 + \lambda_1^{(2)}x_Q)x_P - (2x_P^2 + \lambda_1^{(2)}x_P) - x_P^2x_Q\Big).
\end{aligned}$$

Thus we have the result. □

Corollary 3.12. *(1) The one-form*

$$\Pi^{(4)}{}_{P_1}^{P_2}(P) := \Sigma^{(4)}(P, P_1)dx - \Sigma^{(4)}(P, P_2)dx$$

is a differential of the third kind, whose only (first-order) poles are $P = P_1$ and $P = P_2$, with residues $+1$ and -1 respectively.

(2) The fundamental differential of the second kind $\Omega^{(4)}(P_1, P_2)$ is given by

$$\begin{aligned}
\Omega^{(4)}(P_1, P_2) &= d_{P_2}\Sigma^{(4)}(P_1, P_2) + \sum_{i=1}^{g}\nu_i^{I;4}(P_1)\otimes\nu_i^{II;4}(P_2) \\
&= \frac{F^{(4)}(P_1, P_2)dx_1\otimes dx_2}{(x_1 - x_2)^2 9y_{7,P_1}y_{8,P_1}y_{7,P_2}y_{8,P_2}},
\end{aligned} \tag{3.14}$$

where $F^{(4)}$ is an element of $R_4 \otimes R_4$.

Proof. Direct computations show it. □

Lemma 3.13. *We have*

$$\lim_{P_1\to\infty}\frac{F^{(4)}(P_1, P_2)}{\phi^{(4)}_{H^1 3}(P_1)(x_1 - x_2)^2} = \phi^{(4)}_{H^1 4}(P_2) = x_{P_2}^2 y_{7,P_2}. \tag{3.15}$$

Proof. A calculation, using the formula for $F^{(4)}$ from 3.12 and the expression for B_4 in the proof of Proposition 3.11, gives the result. □

Hereafter for $P = (x, y_7, y_8)$, we expand $h_i(P) = 3y_7y_8\nu_i^{II;4}(P)/dx$, $1 \le i \le g$, in the monomial representatives for elements of R_4 to avoid carrying holomorphic one-forms.

For later convenience we introduce the notation:

$$\begin{aligned}\Omega^{(4)}{}^{P_1,P_2}_{Q_1,Q_2} &:= \int_{P_2}^{P_1}\int_{Q_2}^{Q_1} \Omega^{(4)}(P,Q) \\ &= \int_{P_2}^{P_1}(\Sigma^{(4)}(P,Q_1) - \Sigma^{(4)}(P,Q_2)) + \sum_{i=1}^{4}\int_{P_2}^{P_1}\nu_i^{I;4}(P)\int_{Q_2}^{Q_1}\nu_i^{II;4}(P).\end{aligned} \tag{3.16}$$

4. SEMIGROUP $H = \langle 6, 13, 14, 15, 16\rangle$

In this section, we investigate a non-singular curve whose Weierstrass semigroup at one point ∞ is the numerical semigroup H_{12} generated by $M_{12} = \{6, 13, 14, 15, 16\}$. Again we follow Pinkham's program, deform the local equations of a smooth affine curve with a given monomial ring at ∞, and use Komeda's result to ensure that there exist a global curve in the deformation space.

4.1. **The curve as a monomial curve.**

The proof of the following Proposition is contributed by Komeda.

Proposition 4.1. *Let B_{12} be the monomial ring which is given by $k[t^a]_{a\in M_{12}}$ for the numerical semigroup H_{12}. For the k-algebra homomorphism,*

$$\varphi_{12} : k[Z] := k[Z_6, Z_{13}, Z_{14}, Z_{15}, Z_{16}] \to k[t^a]_{a\in M_{12}},\ Z_a \mapsto t^a,$$

where Z_a has weight $a = 6, 13, 14, 15, 16$, the kernel of φ_{12} is generated by the following relations $f^{(Z)}_{12,b}$ $(b = 1, \cdots, 9)$,

$$\begin{array}{lll} f^{(Z)}_{12,1} = Z_{13}^2 - Z_6^2Z_{14}, & f^{(Z)}_{12,2} = Z_{13}Z_{14} - Z_6^2Z_{15}, & f^{(Z)}_{12,3} = Z_{14}^2 - Z_{13}Z_{15}, \\ f^{(Z)}_{12,4} = Z_{14}^2 - Z_6^2Z_{16}, & f^{(Z)}_{12,5} = Z_{13}Z_{16} - Z_{14}Z_{15}, & f^{(Z)}_{12,6} = Z_{15}^2 - Z_6^5, \\ f^{(Z)}_{12,7} = Z_{14}Z_{16} - Z_6^5, & f^{(Z)}_{12,8} = Z_{15}Z_{16} - Z_6^3Z_{13}, & f^{(Z)}_{12,9} = Z_{16}^2 - Z_6^3Z_{14}. \end{array} \tag{4.1}$$

Proof. We set $a_1 = 6$, $a_2 = 13$, $a_3 = 14$, $a_4 = 15$ and $a_5 = 16$. Let $\varphi_{12} : k[Z_6, Z_{13}, Z_{14}, Z_{15}, Z_{16}] \longrightarrow k[H_{12}] = k[t^h]_{h\in H_{12}}$ be the k-algebra homomorphism which sends Z_i to t^{a_i} for each i. Then the ideal $I = \ker\varphi$ is generated by

$$f^{(Z)}_{12,1} = Z_{13}^2 - Z_6^2Z_{14},\ f^{(Z)}_{12,2} = Z_{13}Z_{14} - Z_6^2Z_{15},\ f^{(Z)}_{12,3} = Z_{14}^2 - Z_{13}Z_{15},$$

$$f^{(Z)}_{12,4} = Z_{14}^2 - Z_6^2Z_{16},\ f^{(Z)}_{12,5} = Z_{13}Z_{16} - Z_{14}Z_{15},\ f^{(Z)}_{12,6} = Z_{15}^2 - Z_6^5,$$

$$f^{(Z)}_{12,7} = Z_{14}Z_{16} - Z_6^5,\ f^{(Z)}_{12,8} = Z_{15}Z_{16} - Z_6^3Z_{13},\ f^{(Z)}_{12,9} = Z_{16}^2 - Z_6^3Z_{14}.$$

We set $\alpha_i = \min\{\alpha \in \mathbb{N}_0 > 0 \mid \alpha a_i = \langle a_1, \dots, a_{i-1}, a_{i+1}, \dots, a_5\rangle\}$. Then it is easy to check that $\alpha_1 = 5$, $\alpha_2 = \alpha_3 = \alpha_4 = \alpha_5 = 2$ and

$$5a_1 = 2a_4,\ 2a_2 = 2a_1 + a_3,\ 2a_4 = 2a_6 + a_5,\ 2a_4 = a_3 + a_5,\ 2a_5 = a_1 + 2a_2.$$

We investigate relations as follows:

$$\beta_i a_i + \beta_j a_j = \gamma_k a_k + \gamma_\ell a_\ell + \gamma_m a_m$$

with $0 < \beta_i < \alpha_i$, $0 < \beta_j < \alpha_j$, $0 < \gamma_k < \alpha_k$, $0 < \gamma_\ell < \alpha_\ell$ and $0 \leqq \gamma_m < \alpha_m$. Then we get the following four relations:

$$a_2 + a_3 = 2a_1 + a_4,\ a_2 + a_4 = 2a_1 + a_5,\ a_2 + a_5 = a_3 + a_4,\ a_4 + a_5 = 3a_1 + a_3.$$

It is easy to check that the polynomials given in the statement belong to I. Let J be the ideal generated by the above nine polynomials. To prove $I = J$ it suffices to show the following: If

$$f = \prod_i Z_{a_i}^{\beta_i} - \prod_i Z_{a_i}^{\gamma_i} \in \ker\varphi \text{ with } \beta_i\gamma_i = 0, \text{ every } i,$$

then $f \in J$.

Case 1. Let $Z_{a_i}^{\beta_i} - Z_{a_j}^{\gamma_j} Z_{a_k}^{\gamma_k} Z_{a_\ell}^{\gamma_\ell} Z_{a_m}^{\gamma_m} \in I$ with $\beta_i \geqq \alpha_i$, $\gamma_j > 0$, $\gamma_k \geqq 0$, $\gamma_\ell \geqq 0$ and $\gamma_m \geqq 0$.

Let $i = 6$. We have

$$f = Z_6^{\beta_1-5}(-f_{12,6}^{(Z)}) + Z_6^{\beta_1-5} Z_{15}^2 - Z_{a_j}^{\gamma_j} M \equiv Z_6^{\beta_1-5} Z_{15}^2 - Z_{a_j}^{\gamma_j} M \bmod J$$

where M is some monomial. Hence, if $j = 4$, then we may decrease the degree of f. If $j = 3$ or 5, using $f_{12,6}^{(Z)} - f_{12,7}^{(Z)}$ we may decrease the degree of f. We may assume that $f = Z_6^{\beta_1} - Z_{13}^{\gamma_2}$ with $\gamma_2 \geqq \alpha_2$. Hence, we have

$$f = Z_6^{\beta_1} - Z_{13}^{\gamma_2-2} f_{12,1}^{(Z)} - Z_{13}^{\gamma_2-2} Z_6^2 Z_{14} \equiv Z_6^{\beta_1} - Z_{13}^{\gamma_2-2} Z_6^2 Z_{14} \bmod J,$$

which implies that we may decrease the degree of f.

Let $i = 2$. We have

$$Z_{13}^{\beta_2} = Z_{13}^{\beta_2-2} f_{12,1}^{(Z)} + Z_{13}^{\beta_2-2} Z_6^2 Z_{14}.$$

Hence, if $j = 1$ or 3, then we may decrease the degree of f. Let $f = Z_{13}^{\beta_2} - Z_{15}^{\gamma_4} Z_{16}^{\gamma_5}$. Using $f_{12,8}^{(Z)} \in J$ we may assume that $\gamma_4 \geqq 2$ or $\gamma_5 \geqq 2$. If $\gamma_5 \geqq 2$, by $f_{12,9}^{(Z)} \in J$ we may decrease the degree of f. Let $\gamma_4 \geqq 2$. Then

$$f = Z_{13}^{\beta_2-2} f_{12,1}^{(Z)} + Z_{13}^{\beta_2-2} Z_6^2 Z_{14} - Z_{15}^{\gamma_4-2}(f_{12,6}^{(Z)} - f_{12,7}^{(Z)}) Z_{16}^{\gamma_5} - Z_{15}^{\gamma_4-2} Z_{14} Z_{16}^{\gamma_5+1}.$$

Hence, we may decrease the degree of f.

Let $i = 14$. Since we obtain

$$Z_{14}^{\beta_3} = Z_{14}^{\beta_3-2} f_{12,4}^{(Z)} + Z_6^2 Z_{14}^{\beta_3-2} Z_{16} = Z_{14}^{\beta_3-2} f_{12,3}^{(Z)} + Z_{13} Z_{14}^{\beta_3-2} Z_{15},$$

which implies that we may decrease the degree of f.

Let $i = 4$. In view of

$$Z_{15}^{\beta_4} = Z_{15}^{\beta_4-2}(f_{12,6}^{(Z)} - f_{12,7}^{(Z)}) + Z_{14} Z_{15}^{\beta_4-2} Z_{16} = Z_{15}^{\beta_4-2} f_{12,6}^{(Z)} + Z_6^5 Z_{15}^{\beta_4-2}$$

we may assume that $f = Z_{15}^{\beta_4} - Z_{13}^{\gamma_2}$. Hence, we get

$$f = Z_{15}^{\beta_4 - 2} f_{12,6}^{(Z)} - Z_{13}^{\gamma_2 - 2} f_{12,1}^{(Z)} + Z_6^5 Z_{15}^{\beta_4 - 2} - Z_6^2 Z_{13}^{\gamma_2 - 2} Z_{14}.$$

We may decrease the degree of f.

Let $i = 16$. Since we have

$$Z_{16}^{\beta_5} = Z_{16}^{\beta_5 - 2} f_{12,9}^{(Z)} + Z_6 Z_{13}^2 Z_{16}^{\beta_5 - 2},$$

we may assume that $f = Z_{16}^{\beta_5} - Z_{14}^{\gamma_3} Z_{15}^{\gamma_4}$. In view of $f_{12,5}^{(Z)} \in J$ we may assume that $\gamma_{14} \geqq 2$ or $\gamma_4 \geqq 2$. By $f_{12,4}^{(Z)}$ and $f_{12,6}^{(Z)} - f_{12,7}^{(Z)} \in J$ we may decrease the degree of f.

It is sufficient to show the following case about the remainder.

Case 2. Let $f = Z_{a_i}^{\beta_i} Z_{a_j}^{\beta_j} - Z_{a_k}^{\gamma_k} Z_{a_\ell}^{\gamma_\ell} Z_{a_m}^{\gamma_m} \in I$ with $\beta_i > 0$, $\beta_j > 0$, $\gamma_k > 0$, $\gamma_\ell > 0$ and $\gamma_m \geqq 0$. By the proof of Case 1 we may assume that $0 < \beta_i < \alpha_i$, $0 < \beta_j < \alpha_j$, $0 < \gamma_k < \alpha_k$, $0 < \gamma_\ell < \alpha_\ell$ and $0 \leqq \gamma_m < \alpha_m$. In this case f is one of the four polynomials $f_{12,2}^{(Z)}$, $f_{12,4}^{(Z)} - f_{12,3}^{(Z)}$, $f_{12,5}^{(Z)}$ and $f_{12,8}^{(Z)}$. ☐

We label $\mathbb{G}_m^{(12)}$ the action of $\mathbb{G}_m$ on $B_{12} \simeq k[Z]/\ker\varphi_{12}$ so that for $g \in \mathbb{G}_m^{(12)}$ $Z_a \mapsto g^a Z_a$ and each a agrees with the weight.

The relations listed in Proposition 4.1 are given by the 2×2 minors of

$$\begin{vmatrix} Z_6^2 & Z_{14} & Z_{16} \\ Z_{14} & Z_{16} & Z_6^3 \end{vmatrix}, \quad \begin{vmatrix} Z_6^2 & Z_{13} & Z_{14} & Z_{15} \\ Z_{13} & Z_{14} & Z_{15} & Z_{16} \end{vmatrix}; \tag{4.2}$$

although $Z_{13}Z_{15} - Z_6^2 Z_{16}$ is not listed, and $f_{12,8}^{(Z)}$ is not one of the minors they are compatible: for example, the given minor follows by combining $f_{12,8}^{(Z)}$ with $f_{12,1}^{(Z)}$ and $f_{12,9}^{(Z)}$. We note that the first matrix is the same as (3.2). Thus, we reprise the notation used in Section 3.

We construct a non-singular curve X_{12} by giving an affine patch, an ideal in the ring $\mathbb{C}[x, y_{13}, y_{14}, y_{15}, y_{16}]$. For any complex numbers b_6 and b_7 distinct from the previous b's, we let

$$\hat{k}_2(x) := (x - b_6)(x - b_7) = x^2 + \hat{\lambda}_1^{(2)} x + \hat{\lambda}_0^{(2)}, \qquad \hat{k}_5(x) := \hat{k}_2(x) k_3(x),$$
$$k_{13}(x) := k_3(x) k_2(x)^2 \hat{k}_2(x)^3, \qquad k_{14}(x) := k_7(x)^2 = k_3(x)^2 k_2(x)^4,$$
$$k_{15}(x) := \hat{k}_5(x)^3, \qquad k_{16}(x) := k_8(x)^2 = k_3(x)^4 k_2(x)^2,$$

Let the prime ideal $\mathcal{P}$ in $\mathbb{C}[x, y_{13}, y_{14}, y_{15}, y_{16}]$ be defined by

$$\mathcal{P} := (f_{12,1}, f_{12,2}, f_{12,3}, f_{12,4}, f_{12,5}, f_{12,6}, f_{12,7}, f_{12,8}, f_{12,9}),$$

$$f_{12,1} := y_{13}^2 - \hat{k}_2(x) y_{14}, \quad f_{12,2} := y_{13} y_{14} - k_2(x) y_{15}, \quad f_{12,3} := \hat{k}_2(x) y_{14}^2 - y_{13} y_{15} k_2(x)$$
$$f_{12,4} := y_{14}^2 - k_2(x) y_{16}, \quad f_{12,5} := y_{13} y_{16} - y_{14} y_{15}, \quad f_{12,6} := y_{15}^2 - \hat{k}_2(x) k_3(x),$$
$$f_{12,7} := y_{14} y_{16} - k_2(x) k_3(x), \quad f_{12,8} := y_{15} y_{16} - k_3(x) y_{13}, \quad f_{12,9} := y_{16}^2 - k_3(x) y_{14},$$

which are the 2×2 minors of

$$\begin{vmatrix} k_2(x) & y_{14} & y_{16} \\ y_{14} & y_{16} & k_3(x) \end{vmatrix}, \quad \begin{vmatrix} \hat{k}_2(x) & y_{13} & y_{14}\hat{k}_2(x) & y_{15} \\ y_{13} & y_{14} & y_{15}k_2(x) & y_{16} \end{vmatrix}; \tag{4.3}$$

again, the minor $y_{13}y_{15} - \hat{k}_2(x)y_{16}$ is not in the list of $f_{i,j}$ and $f_{12,8}$ is not a minor, but they are compatible–the minor follows by combining $f_{12,8}$ with $f_{12,1}$ and $f_{12,9}$. Here the first matrix is the same as (3.4) and the latter is related to the double covering of X_4. In other words, we essentially identify y_a and y_{2a} for $a = 7, 8$. There is only one point at infinity in this affine model because, as is clear from the equations, x approaches ∞ if and only if each y_b ($b = 13, 14, 15, 16$) approaches ∞. We define the $\mathbb{G}_m^{(12)}$ action on x and y_a by $g_m^{-6}x$ and $g_m^{-a}y_a$ ($a = 13, 14, 15, 16$) near $\infty \in X_{12}$.

Corresponding to Proposition 4.1, we will consider a commutative ring,

$$R_{12} = \mathbb{C}[x, y_{13}, y_{14}, y_{15}, y_{16}]/\mathcal{P}.$$

Proposition 4.2. *There is a ring homomorphism $R_4 \to R_{12}$.*

Following Nagata's Jacobi criterion [Mat, Theorem 30.10], we show that $\mathrm{Spec} R_{12}$ is non-singular.

Proposition 4.3. *For every $(x, y_{13}, y_{14}, y_{15}, y_{16})$ which is a zero of every $(f_{12,a})_{a=1,\cdots,9}$, we have*

$$\mathrm{rank}\, \mathcal{U}_{12} = 4, \quad \mathcal{U}_{12} := \left(\left(\tfrac{\partial}{\partial x} f_{12,a} \quad \tfrac{\partial}{\partial y_{13}} f_{12,a} \quad \tfrac{\partial}{\partial y_{14}} f_{12,a} \quad \tfrac{\partial}{\partial y_{15}} f_{12,a} \quad \tfrac{\partial}{\partial y_{16}} f_{12,a}\right)_{a=1,\cdots,9}\right).$$

Proof. $\mathcal{U}_{12}$ is

$$\begin{pmatrix} -\hat{k}_2' y_{14} & 2y_{13} & -\hat{k}_2 & & \\ -k_2' y_{15} & y_{14} & y_{13} & -k_2 & \\ \hat{k}_2' y_{14}^2 - y_{13}y_{15}k_2' & -y_{15}k_2 & 2\hat{k}_2 y_{14} & -y_{13}k_2 & \\ -k_2' y_{16} & & 2y_{14} & & k_2 \\ & y_{16} & -y_{15} & -y_{14} & y_{13} \\ -(\hat{k}_2 k_3)' & & & 2y_{15} & \\ -(k_2k_3)' & & y_{16} & & y_{14} \\ -k_3' y_{13} & -k_3 & & y_{16} & y_{15} \\ -k_3' y_{14} & & -k_3 & & 2y_{16} \end{pmatrix}.$$

When $x = b_1$ as a zero of $k_3(x)$, every y_a vanishes and thus

$$\begin{pmatrix} & & -\hat{k}_2(b_1) & & \\ & & & -k_2(b_1) & \\ & & & & \\ & & & & \\ -\hat{k}_2(b_1) & & & 2y_{15} & \\ -k_2(b_1)(b_1-b_2)(b_1-b_3) & & & & \\ & & & & \\ & & & & \end{pmatrix}$$

is evidently a matrix of rank 4. When $x = b_4$ as a zero of $k_2(x)$, y_{13}, y_{14} and y_{16} vanish and $\mathcal{U}_{12}$ is equal to

$$\begin{pmatrix}
 & & -\hat{k}_2(b_4) & & \\
-(b_4-b_5)y_{15} & & & & \\
y_{15} & & & & \\
 & & & & \\
 & & & & \\
 & & -y_{15} & & \\
-(\hat{k}_2k_3)'(b_1) & & & 2y_{15} & \\
-(b_4-b_5)k_3(b_4) & & & & \\
 & -k_3(b_4) & & & y_{15}
\end{pmatrix}$$

whose rank is obviously 4. When $x = b_6$ as a zero of $\hat{k}_2(x)$, y_{13} and y_{15} vanish and thus the matrix is

$$\begin{pmatrix}
-(b_6-b_7)y_{14} & & & & \\
 & y_{14} & & -k_2(b_6) & \\
(b_6-b_7)y_{14}^2 & & & & \\
-k_2(b_6)'y_{16} & & 2y_{14} & & k_2(b_6) \\
 & y_{16} & & y_{14} & \\
-(b_6-b_7)k_3(b_6) & & & & \\
-(k_2k_3)'(b_6) & & y_{16} & & y_{14} \\
 & -k_3 & & y_{16} & \\
-k_3'(b_6)y_{14} & & -k_3(b_6) & & 2y_{16}
\end{pmatrix}.$$

Since $\dfrac{y_{14}}{k_3(b_6)} = \dfrac{y_{16}}{y_{14}}$ and $\dfrac{y_{16}}{y_{14}} = \dfrac{k_3(b_6)}{y_{16}} = \dfrac{y_{14}}{k_2(b_6)}$, the rank is 4. When x differs from b_1, b_2, $\cdots$, and b_7, we compute

$$\begin{pmatrix}
1 & & & & & & & & \\
 & 1 & & -y_{15}/y_{16} & -y_{14}/y_{16} & & & & \\
y_{14} & -y_{13} & 1 & & & & & & \\
 & & & 1 & & & & & \\
 & & & & 1 & & & & \\
-k_3/y_{14} & & & & y_{15}/y_{14} & 1 & -\hat{k}_2/y_{13} & & \\
 & -k_3/y_{15} & & & & & 1 & k_2/y_{13} & \\
 & & & & & & & 1 & \\
 & & & & -y_{16}/y_{13} & & & -y_{14}/y_{13} & 1
\end{pmatrix}\mathcal{U}_{12},$$

which is equal to a matrix of rank 4,

$$\begin{pmatrix} -\hat{k}_2' y_{14} & 2y_{13} & -\hat{k}_2 & & \\ & & & & \\ -k_2' y_{16} & & 2y_{14} & & k_2 \\ & y_{16} & -y_{15} & -y_{14} & y_{13} \\ & & & & \\ -k_3' y_{13} & -k_3 & & y_{16} & y_{15} \end{pmatrix}.$$

$\square$

There is a Riemann surface, which we denote X_{12}, obtained from the affine smooth curve given by $\mathrm{Spec}(R_{12})$ by adding one point ∞ (see Appendix B).

In the local ring of the place ∞, we express x and y's as

$$y_a = \frac{1}{t^{12+a}}, \quad (a = 1, 2, 3, 4), \quad x = \frac{1}{t^6},$$

using the local parameter t at ∞.

For later use, we introduce the following ring,

$$\hat{R}_{12} := \mathbb{C}[x, w_3, w_2, \hat{w}_2]/(w_3^6 - k_3(x), w_2^6 - k_2(x), \hat{w}_2^6 - \hat{k}_2(x)),$$

and then we have a natural ring homomorphism $i_{12} : R_{12} \to \hat{R}_{12}$. Since in $\hat{R}_{12}$, we have

$$w_3^6 = k_3(x), \quad w_2^6 = k_2(x), \quad \hat{w}_2^6 = \hat{k}_2(x), \tag{4.4}$$

in $i_{12}(R_{12})$, the following holds

$$y_{13} = w_3 w_2^2 \hat{w}_2^3, \qquad y_{14} = w_3^2 w_2^4, \qquad y_{15} = w_3^3 \hat{w}_2^3, \qquad y_{16} = w_3^4 w_2^2. \tag{4.5}$$

This gives a projection

$$\mathrm{Spec}\ \hat{R}_{12} \to \mathrm{Spec}\ R_{12}. \tag{4.6}$$

The curve X_{12} has a cyclic action of a primitive sixth root of unity ζ_6:

$$\hat{\zeta}_6(x, y_{13}, y_{14}, y_{15}, y_{16}) = (x, \zeta_6 y_{13}, \zeta_6^2 y_{14}, \zeta_6^3 y_{15}, \zeta_6^4 y_{16}).$$

Using the above expression of w's, it is obvious that the prime ideal $\mathcal{P}$ is stable for the cyclic action.

4.2. **The Weierstrass gaps and holomorphic one forms.** The Weierstrass gap sequence of R_{12} at ∞ is given by the following table: in particular, the semigroup $H(X_{12}, \infty)$ equals H_{12}.

Table 2

0	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23
1	-	-	-	-	-	x	-	-	-	-	-	x^2	y_{13}	y_{14}	y_{15}	y_{16}	-	x^3	xy_{13}	xy_{14}	xy_{15}	xy_{16}	-

We introduce a notation for the monomials in R_{12} whose orders of pole at ∞ correspond to the non-gaps:

$$\phi_0^{(12)} = 1, \quad \phi_1^{(12)} = x, \quad \phi_2^{(12)} = x^2, \quad \phi_{a+2}^{(12)} = y_a, \quad \phi_7^{(12)} = x^3, \quad \phi_{a+7}^{(12)} = xy_a,$$

$$\phi_{6i-12}^{(12)} = x^i, \quad \phi_{6i-12+a}^{(12)} = x^{i-1}y_a, \quad \phi_{6i-7}^{(12)} = x^{i-3}y_{13}y_{16}, \quad (a = 1, 2, 3, 4, i = 4, 5, 6, \cdots).$$

We define the "weight"

$$N^{(12)}(n) = -\mathrm{wt}(\phi_n^{(12)}),$$

where wt() is the order of pole at ∞, consistent with the $\mathbb{G}_m^{(12)}$ action.

The related Young diagram $\mathfrak{Y}_{12}$ of M_{12} is given by

$$\Lambda_i = N^{(12)}(g) - N^{(12)}(i-1) - 13 + i,$$

which is equal to $\alpha_{12-i}(L(H_{12}))$ in (2.1).



When b's are distinct, a basis of the holomorphic one-form is given by $\nu_i^{I;12}$, $i = 1, 2, \cdots, 12$,

$$\nu_{[13-i]}^{I;12} := \nu_i^{I;12} := \frac{\phi_{i-1}^{(12)}dx}{6y_{13}y_{16}} = \frac{\phi_{i-1}^{(12)}dx}{6y_{14}y_{15}},$$

or

$$
\begin{aligned}
&\nu_1^{I;12}=\frac{dx}{6y_{13}y_{16}}, \quad \nu_2^{I;12}=\frac{xdx}{6y_{13}y_{16}}, \quad \nu_3^{I;12}=\frac{x^2dx}{6y_{13}y_{16}}, \quad \nu_4^{I;12}=\frac{dx}{6y_{16}},\\
&\nu_5^{I;12}=\frac{dx}{6y_{15}}, \quad \nu_6^{I;12}=\frac{dx}{6y_{14}}, \quad \nu_7^{I;12}=\frac{dx}{6y_{13}y_{16}}, \quad \nu_8^{I;12}=\frac{x^3dx}{6y_{13}y_{16}},\\
&\nu_9^{I;12}=\frac{xdx}{6y_{16}}, \quad \nu_{10}^{I;12}=\frac{xdx}{6y_{15}}, \quad \nu_{11}^{I;12}=\frac{xdx}{6y_{14}}, \quad \nu_{12}^{I;12}=\frac{xdx}{6y_{13}}.
\end{aligned}
\tag{4.7}
$$

and its Abelian integral for a point $P \in X_{12}$ is defined;

$$\hat{u}(P)=\int_\infty^P \nu^{I;12} \in \mathbb{C}^{12},$$

for k points $P_1, P_2, \cdots, P_k \in X_{12}$, the Abelian map $\hat{u}(P_1, \cdots, P_k)$ is defined by $\sum_{i=1}^k \hat{u}(P_i)$. We also choose a basis $\alpha_i^{(12)}, {\beta^{(12)}}_j$ $(1 \leqq i, j \leqq 12)$ of $H_1(X_{12}, \mathbb{Z})$ such that the intersection numbers are $\alpha_i^{(12)} \cdot \alpha_j^{(12)} = {\beta^{(12)}}_i \cdot {\beta^{(12)}}_j = 0$ and $\alpha_i^{(12)} \cdot {\beta^{(12)}}_j = \delta_{ij}$. Let the period matrices be denoted by

$$\left[\, \omega^{(12)\prime} \ \omega^{(12)\prime\prime} \right] := \frac{1}{2}\left[\int_{\alpha_i^{(12)}} \nu_j^{I;12} \quad \int_{{\beta^{(12)}}_i} \nu_j^{I;12} \right]_{i,j=1,2,\cdots,12}, \tag{4.8}$$

Then we define the Jacobian $\mathcal{J}_{12}$ by

$$\kappa : \mathbb{C}^{12} \to \mathcal{J}_{12} = \mathbb{C}^{12}/\Lambda_{12},$$

where Λ_{12} is generated by $\omega^{(12)\prime}$ and $\omega^{(12)\prime\prime}$.

The Abel map of for $(P_1, \cdots, P_k) \in S^k X_{12}$ to $\mathbb{C}^{12}$ is also expressed by

$$\mathsf{t}^{[k]}_{23-N^{(12)}(12-i)} := u_i^{[k]} := \hat{u}(P_1, \cdots, P_k) \in \mathbb{C}^{12}.$$

Further we define the subvariety ${\mathcal{W}^{(12)}}_k$ by

$${\mathcal{W}^{(12)}}_k := \hat{u}(S^k X_{12})/\Lambda_{12}.$$

By letting

$$\mathsf{t}_{23-N^{(12)}(12-i)} := u_i := \mathsf{t}^{[12]}_{23-N^{(12)}(12-i)} \equiv u_i^{[12]}, \tag{4.9}$$

the weights are given by

$$\{\mathrm{wt}(\mathsf{t}_i)\} = \{1, 2, 3, 4, 5, 7, 8, 9.10, 11, 17, 23\}.$$

Here we use the convention that for $P_a \in X_{12}$, P_a is expressed by $(x_a, y_{13,a}, y_{14,a}, y_{15,a}, y_{16,a})$ or $(x_{P_a}, y_{13,P_a}, y_{14,P_a}, y_{15,P_a}, y_{16,P_a})$ and $w_{b,a}$ is also w_b associated with P_a.

By letting $u := \hat{u}(P_1, \cdots, P_{12})$, we have

$$\begin{pmatrix} \partial/\partial u_1 \\ \partial/\partial u_2 \\ \vdots \\ \partial/\partial u_{11} \\ \partial/\partial u_{12} \end{pmatrix} = \Psi_{12}^{(12)-1} \begin{pmatrix} 6y_{13,1}y_{14,1}\partial/\partial x_1 \\ 6y_{13,2}y_{14,2}\partial/\partial x_2 \\ \vdots \\ 6y_{13,11}y_{14,11}\partial/\partial x_{11} \\ 6y_{13,12}y_{14,12}\partial/\partial x_{12} \end{pmatrix},$$

where

$$\Psi_{12}^{(12)} := \begin{pmatrix} \phi_0^{(12)}(P_1) & \phi_1^{(12)}(P_1) & \phi_2^{(12)}(P_1) & \cdots & \phi_{11}^{(12)}(P_1) \\ \phi_0^{(12)}(P_2) & \phi_1^{(12)}(P_2) & \phi_2^{(12)}(P_2) & \cdots & \phi_{11}^{(12)}(P_2) \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ \phi_0^{(12)}(P_{11}) & \phi_1^{(12)}(P_{11}) & \phi_2^{(12)}(P_{11}) & \cdots & \phi_{11}^{(12)}(P_{11}) \\ \phi_0^{(12)}(P_{12}) & \phi_1^{(12)}(P_{12}) & \phi_2^{(12)}(P_{12}) & \cdots & \phi_{11}^{(12)}(P_{12}) \end{pmatrix}. \tag{4.10}$$

In particular, for any 12-tuple of numbers $(\varepsilon_i)_{i=1,\dots,12}$, we have

$$\sum_{i=1} \epsilon_i \frac{\partial}{\partial u_i} = |\Psi_{12}^{(12)-1}| \begin{vmatrix} \phi_0^{(12)}(P_1) & \phi_1^{(12)}(P_1) & \phi_2^{(12)}(P_1) & \cdots & \phi_{11}^{(12)}(P_1) & 3y_{13,1}y_{16,1}\partial/\partial x_1 \\ \phi_0^{(12)}(P_2) & \phi_1^{(12)}(P_2) & \phi_2^{(12)}(P_2) & \cdots & \phi_{11}^{(12)}(P_2) & 3y_{13,2}y_{16,2}\partial/\partial x_2 \\ \vdots & \vdots & \vdots & \ddots & \vdots & \\ \phi_0^{(12)}(P_{11}) & \phi_1^{(12)}(P_{11}) & \phi_2^{(12)}(P_{11}) & \cdots & \phi_{11}^{(12)}(P_{11}) & 3y_{13,3}y_{16,3}\partial/\partial x_3 \\ \phi_0^{(12)}(P_{12}) & \phi_1^{(12)}(P_{12}) & \phi_2^{(12)}(P_{12}) & \cdots & \phi_{11}^{(12)}(P_{12}) & 3y_{13,4}y_{16,4}\partial/\partial x_4 \\ \epsilon_1 & \epsilon_2 & \epsilon_3 & \cdots & \epsilon_{12} & \end{vmatrix}.$$

We express the change of variables:

$$\sum_{i,j=1}^{12} \phi_{i-1}^{(12)}(P_1)\phi_{j-1}^{(12)}(P_2) \frac{\partial^2}{\partial \hat{u}_i(P_1)\partial \hat{u}_j(P_2)} = 9y_{13,1}y_{16,1}y_{13,2}y_{16,2}\frac{\partial^2}{\partial x_1 \partial x_2}. \tag{4.11}$$

4.3. **Differentials of the second and the third kinds.** We also give an algebraic representation of the fundamental normalized differential of the second kind $\Omega^{(12)}(P_1, P_2)$ on $X_{12} \times X_{12}$, whose defining properties are the same as those given for $\Omega^{(4)}(P_1, P_2)$ on $X_4 \times X_4$.

Proposition 4.4. *Let $\Sigma^{(12)}\big(P_1, P_2\big)$ be the following form,*

(4.12)

$$\Sigma^{(12)}(P_1, P_2) := \frac{y_{13,1}y_{16,1} + y_{13,2}y_{16,1} + y_{13,1}y_{16,2} + y_{14,2}y_{15,1} + y_{14,1}y_{15,2} + y_{14,2}y_{15,2}}{6(x_1 - x_2)y_{13,1}y_{16,1}} dx_1.$$

Then $\Sigma^{(12)}(P, Q)$ has the following properties

1. *$\Sigma^{(12)}(P, Q)$ as a function of P is singular at $Q = (x_Q, y_{13,Q}, y_{14,Q}, y_{15,Q}, y_{16,Q})$ and ∞ and vanishes at $\hat{\zeta}_6(Q) = (x_Q, \zeta_6 y_{13,Q}, \zeta_6^2 y_{14,Q}, \zeta_6^3 y_{15,Q}, \zeta_6^4 y_{16,Q})$.*
2. *$\Sigma^{(12)}(P, Q)$ as a function of Q has singular at P and at ∞.*

Proof. The result can be checked more easily in terms of the functions w's in the ring $\hat{R}_{12}$, cf. (4.4); we express $\Sigma^{(12)}(P_1, P_2)$ in term of w's,

$$\begin{aligned}\Sigma^{(12)}(P_1, P_2) = &\frac{w_{3,1} w_{2,1}^2 \hat{w}_{2,1}^3 w_{3,1}^4 w_{2,1}^3 + w_{3,2} w_{2,2}^2 \hat{w}_{2,2}^3 w_{3,1}^4 w_{2,1}^3 + w_{3,1} w_{2,1}^2 \hat{w}_{2,1}^3 w_{3,2}^4 w_{2,2}^3}{6(x_1 - x_2) y_{13,1} y_{16,1}} dx_1 \\ &+ \frac{w_{3,1}^2 w_{2,1}^4 w_{3,1}^4 \hat{w}_{2,1}^3 + w_{3,1}^2 w_{2,1}^4 w_{3,2}^4 \hat{w}_{2,2}^3 + w_{3,2}^2 w_{2,2}^4 w_{3,1}^4 \hat{w}_{2,1}^3}{6(x_1 - x_2) y_{13,1} y_{16,1}} dx_1.\end{aligned} \tag{4.13}$$

When the x-coordinates of P_1 and P_2 differ, it obvious that it does not diverges. When the x-coordinates coincides but P_1 and P_2 differ, some of $w_{3,1}$, $w_{2,1}$, and $\hat{w}_{3,1}$ differ from $w_{3,2}$, $w_{2,2}$, and $\hat{w}_{3,2}$. Then the numerator vanishes to order higher than two when expanded in the local parameter. $\Sigma^{(12)}(P_1, P_2)$ diverges only at infinity. ☐

The following holds for smooth X_{12};

Proposition 4.5. *There exist differentials* $\nu_j^{II;12} = \nu_j^{II;12}(x, y)$ $(j = 1, 2, \cdots, 12)$ *of the second kind such that they have their only pole at* ∞ *and satisfy the relation,*

$$\begin{aligned} &d_Q \Sigma^{(12)}\big(P, Q\big) - d_P \Sigma^{(12)}\big(Q, P\big) \\ &\quad = \sum_{i=1}^{12} \Big(\nu_i^{I;12}(Q) \otimes \nu_i^{II;12}(P) - \nu_i^{I;12}(P) \otimes \nu_i^{II;12}(Q) \Big), \end{aligned} \tag{4.14}$$

where

$$d_Q \Sigma^{(12)}\big(P, Q\big) := dx_P \otimes dx_Q \frac{\partial}{\partial x_Q} \Sigma^{(12)}. \tag{4.15}$$

The set of differentials $\{\nu_1^{II;12}, \nu_2^{II;12}, \cdots, \nu_{12}^{II;12}\}$ *is determined modulo the* $\mathbb{C}$*-linear space spanned by* $\langle \nu_j^{I;12} \rangle_{j=1,\ldots,12}$.

Proposition 4.6. *The differentials of the second kind are given by,*

$$\begin{aligned}\nu_1^{II;12} = -\Big(&23x^5 + (19\lambda_1^{(2)} + 18\lambda_1^{(3)} + 20\hat{\lambda}_1^{(2)})x^4 + (14\lambda_1^{(3)}\lambda_1^{(2)} + 15\lambda_1^{(3)}\hat{\lambda}_1^{(2)} + 16\lambda_1^{(2)}\hat{\lambda}_1^{(2)} \\ &+ 13\lambda_2^{(3)} + 15\lambda_2^{(2)} + 17\hat{\lambda}_2^{(2)})x^3 \\ &+ (10\lambda_1^{(3)}\lambda_2^{(2)} + 9\lambda_2^{(3)}\lambda_1^{(2)} + 10\lambda_2^{(3)}\hat{\lambda}_1^{(2)} + 12\lambda_2^{(2)}\hat{\lambda}_1^{(2)} + 12\lambda_1^{(3)}\hat{\lambda}_2^{(2)} + 13\lambda_1^{(2)}\hat{\lambda}_2^{(2)} \\ &+ 8\lambda_3^{(3)} + 11\lambda_1^{(3)}\lambda_1^{(2)}\hat{\lambda}_1^{(2)})x^2 \\ &+ (4\lambda_3^{(3)}\lambda_1^{(2)} + 5\lambda_2^{(3)}\lambda_2^{(2)} + 5\lambda_3^{(3)}\hat{\lambda}_1^{(2)} + 7\lambda_2^{(3)}\hat{\lambda}_2^{(2)} + 9\lambda_2^{(2)}\hat{\lambda}_2^{(2)} + 6\lambda_2^{(3)}\lambda_1^{(2)}\hat{\lambda}_1^{(2)} \\ &+ 8\lambda_1^{(3)}\lambda_1^{(2)}\hat{\lambda}_2^{(2)} + 7\lambda_1^{(3)}\lambda_2^{(2)}\hat{\lambda}_1^{(2)})x \\ &+ 3\lambda_2^{(3)}\lambda_1^{(2)}\hat{\lambda}_2^{(2)} + 4\lambda_1^{(3)}\lambda_2^{(2)}\hat{\lambda}_2^{(2)} + 2\lambda_2^{(3)}\lambda_2^{(2)}\hat{\lambda}_1^{(2)} + \lambda_3^{(3)}\lambda_1^{(2)}\hat{\lambda}_1^{(2)} + 2\lambda_3^{(3)}\hat{\lambda}_2^{(2)}\Big) dx/6y_{13},\end{aligned}$$

$$\begin{aligned}
\nu_2^{II;12} = -\Big(&17x^4 + (13\lambda_1^{(2)} + 12\lambda_1^{(3)} + 14\hat{\lambda}_1^{(2)})x^3 \\
&+ (10\lambda_1^{(2)}\hat{\lambda}_1^{(2)} + 9\lambda_1^{(3)}\hat{\lambda}_1^{(2)} + 8\lambda_1^{(3)}\lambda_1^{(2)} + 11\hat{\lambda}_2^{(2)} + 9\lambda_2^{(2)} + 7\lambda_2^{(3)})x^2 \\
&+ (7\lambda_1^{(2)}\hat{\lambda}_2^{(2)} + 6\lambda_1^{(3)}\hat{\lambda}_2^{(2)} + 4\lambda_1^{(3)}\lambda_2^{(2)} + 6\lambda_2^{(2)}\hat{\lambda}_1^{(2)} + 4\lambda_2^{(3)}\hat{\lambda}_1^{(2)} + 3\lambda_2^{(3)}\lambda_1^{(2)} \\
&\quad + 2\lambda_3^{(3)} + 5\lambda_1^{(3)}\lambda_1^{(2)}\hat{\lambda}_1^{(2)})x \\
&+ \lambda_1^{(3)}\lambda_2^{(2)}\hat{\lambda}_1^{(2)} + 2\lambda_1^{(3)}\lambda_1^{(2)}\hat{\lambda}_2^{(2)} + 3\lambda_2^{(2)}\hat{\lambda}_2^{(2)} + \lambda_2^{(3)}\hat{\lambda}_2^{(2)}\Big)dx/6y_{13},
\end{aligned}$$

$$\begin{aligned}
\nu_3^{II;12} =&\Big(11x^3 + (6\lambda_1^{(3)} + 7\lambda_1^{(2)} + 8\hat{\lambda}_1^{(2)})x^2 + (4\lambda_1^{(2)}\hat{\lambda}_1^{(2)} + 3\lambda_1^{(3)}\hat{\lambda}_1^{(2)} + 2\lambda_1^{(3)}\lambda_1^{(2)} \\
&\quad + \lambda_2^{(3)} + 3\lambda_2^{(2)} + 5\hat{\lambda}_2^{(2)})x - 3\lambda_2^{(3)}\lambda_1^{(2)} + \lambda_1^{(2)}\hat{\lambda}_2^{(2)}\Big)dx/6y_{13}, \\
\nu_4^{II;12} =& -\Big((8x^3 + (4\lambda_1^{(2)} + 6\lambda_1^{(3)})x^2 + (2\lambda_1^{(3)}\lambda_1^{(2)}4\lambda_2^{(3)})x + 2\lambda_3^{(3)})\Big)dx/6y_{14},
\end{aligned}$$

$$\begin{aligned}
\nu_5^{II;12} &= -\Big(9x^3 + (6\hat{\lambda}_1^{(2)} + 6\lambda_1^{(3)})x^2 + (3\lambda_1^{(3)}\hat{\lambda}_1^{(2)} + 3\hat{\lambda}_2^{(2)} + 3\lambda_2^{(3)})x\Big)dx/6y_{15}, \\
\nu_6^{II;12} &= -\Big(10x^3 + (8\lambda_1^{(2)} + 6\lambda_1^{(3)})x^2 + (4\lambda_1^{(3)}\lambda_1^{(2)} + 2\lambda_2^{(3)} + 6\lambda_2^{(2)})x + 2\lambda_1^{(3)}\lambda_2^{(2)}\Big)dx/6y_{16},
\end{aligned}$$

$$\begin{aligned}
\nu_7^{II;12} &= -\Big(7x^5 + (6\lambda_1^{(3)} + 5\lambda_1^{(2)} + 4\hat{\lambda}_1^{(2)})x^4 \\
&+ (3\lambda_2^{(2)} + 3\lambda_1^{(3)}\hat{\lambda}_1^{(2)} - \hat{\lambda}_2^{(2)} + 5\lambda_2^{(3)} + 2\lambda_1^{(2)}\hat{\lambda}_1^{(2)} + 4\lambda_1^{(3)}\lambda_1^{(2)})x^3 \\
&+ (\lambda_1^{(3)}\lambda_1^{(2)}\hat{\lambda}_1^{(2)} + 2\lambda_2^{(3)}\hat{\lambda}_1^{(2)} + 3\lambda_2^{(3)}\lambda_1^{(2)} + 2\lambda_1^{(3)}\lambda_2^{(2)} + 4\lambda_3^{(3)})x^2 \\
&+ (-2\lambda_3^{(3)}\lambda_1^{(2)} - \lambda_2^{(3)}\lambda_2^{(2)} - \lambda_3^{(3)}\hat{\lambda}_1^{(2)})\Big)dx/6y_{13}y_{16}, \\
\nu_8^{II;12} &= -\Big(5x^2 + (2\hat{\lambda}_1^{(2)} + \lambda_1^{(2)})x\Big)/6y_{13}, \\
\nu_9^{II;12} &= -2x^2dx/6y_{14}, \qquad \nu_{10}^{II;12} = -3x^2dx/6y_{15}, \qquad \nu_{11}^{II;12} = -(2\lambda_1^{(2)}x + 4x^2)dx/6y_{16}, \\
\nu_{12}^{II;12} &= -x^4/6y_{13}y_{16}.
\end{aligned}$$

Proof. By denoting the numerator in (4.12) and (4.13) by denoted by

$$\begin{aligned}
A(P_1, P_2) :=& \; y_{13,1}y_{16,1} + y_{13,2}y_{16,1} + y_{13,1}y_{16,2} + y_{14,2}y_{15,1} + y_{14,1}y_{15,2} + y_{14,2}y_{15,2} \\
=& \; w_{3,1}w_{2,1}^2\hat{w}_{2,1}^3w_{3,1}^4w_{2,1}^2 + w_{3,2}w_{2,2}^2\hat{w}_{2,2}^3w_{3,1}^4w_{2,1}^2 + w_{3,1}w_{2,1}^2\hat{w}_{2,1}^3w_{3,2}^4w_{2,2}^2 \\
&\quad + w_{3,1}w_{2,1}^4w_{3,1}^4\hat{w}_{2,1}^3 + w_{3,1}w_{2,1}^4w_{3,2}^4\hat{w}_{2,2}^3 + w_{3,2}w_{2,2}^4w_{3,1}^4\hat{w}_{2,1}^3 \\
=& \; w_{3,1}^5w_{2,1}^4\hat{w}_{2,1}^3 + w_{3,2}w_{2,2}^2\hat{w}_{2,2}^3w_{3,1}^4w_{2,1}^2 + w_{3,1}w_{2,1}^2\hat{w}_{2,1}^3w_{3,2}^4w_{2,2}^2 \\
&\quad + w_{3,2}^2w_{2,2}^3w_{3,1}^3\hat{w}_{2,1}^3 + w_{3,1}^2w_{2,1}^3w_{3,2}^3\hat{w}_{2,2}^3 + w_{3,2}^5w_{2,2}^4\hat{w}_{2,1}^3,
\end{aligned}$$

we have the equality,

$$
\begin{aligned}
&\frac{\partial \Sigma^{(12)}(P_1,P_2)}{\partial x_2} - \frac{\partial \Sigma^{(12)}(P_2,P_1)}{\partial x_1} \\
&\qquad = \frac{(y_{13,2}y_{16,2}A(P_1,P_2) - y_{13,1}y_{16,1}A(P_2,P_1))/(x_1-x_2)}{(x_1-x_2)y_{13,1}y_{16,1}y_{13,2}y_{16,2}} \\
&\qquad + \frac{y_{13,1}y_{16,1}\frac{\partial A(P_1,P_2)}{\partial x_2} + y_{13,2}y_{16,2}\frac{\partial A(P_2,P_1)}{\partial x_1}}{(x_1-x_2)y_{13,1}y_{16,1}y_{13,2}y_{16,2}}.
\end{aligned} \tag{4.16}
$$

We must evaluate the numerator of (4.16) which is denoted by $B(P_1,P_2)$. The former terms are written by,

$$
\begin{aligned}
&y_{13,2}y_{16,2}A(P_1,P_2) - y_{13,1}y_{16,1}A(P_2,P_1) \\
&= w_{3,2}^6 w_{2,2}^6 \hat{w}_{2,2}^6 w_{3,1}^4 w_{2,1}^2 + w_{3,1}w_{2,1}^2\hat{w}_{2,1}^3 w_{3,2}^9 w_{2,2}^6 {w'_{2,2}}^3 + w_{3,2}^7 w_{2,2}^8 \hat{w}_{2,2}^3 w_{3,1}^4 \hat{w}_{2,1}^3 \\
&+ w_{3,1}w_{2,1}^4 w_{3,2}^8 w_{2,2}^4 \hat{w}_{2,2}^6 + w_{3,2}^{10} w_{2,2}^8 \hat{w}_{2,1}^6 - (P_1 \leftrightarrow P_2) \\
&= y_{16,1}k_{3,2}k_{2,2}\hat{k}_{2,2} + y_{13,1}y_{15,2}k_{3,2}k_{2,2} + y_{13,2}y_{15,1}k_{3,2}k_{2,2} + y_{14,1}y_{14,2}k_{3,2}\hat{k}_{2,2} \\
&+ y_{16,2}k_{3,2}k_{2,2}\hat{k}_{2,1} - (P_1 \leftrightarrow P_2) \\
&= y_{16,1}(k_{3,2}k_{2,2}\hat{k}_{2,2} - k_{3,1}k_{2,1}\hat{k}_{2,1}) + y_{16,2}(k_{3,2}k_{2,2}\hat{k}_{2,2} - k_{3,1}k_{2,1}\hat{k}_{2,1}) \\
&+ (y_{13,1}y_{15,2} + y_{13,2}y_{15,1})(k_{3,2}k_{2,2} - k_{3,1}k_{2,1}) + y_{14,1}y_{14,2}(k_{3,2}\hat{k}_{2,2} - k_{3,1}\hat{k}_{2,1}).
\end{aligned}
$$

Here $k_{a,b} := k_a(x_b)$, and $\hat{k}_{2,b} := \hat{k}_2(x_b)$ for $a = 2,3$ and $b = 1,2$ and $k' := dk/dx$. We reinforce that these are expressions in the affine coordinates x's and y's, in other words the w's in these expressions, which are algebraic over the ring R_{12}, appear only within algebraic combinations which belong to R_{12}.

The differential terms of (4.16) are given as follows:

$$
\begin{aligned}
&6y_{13,2}y_{16,2}\frac{\partial A(P_2,P_1)}{\partial x_2} = w_{3,2}^5 w_{2,2}^4 {w'_{2,2}}^3 \Big(y_{16,1}\Big(\frac{1}{w_{3,2}^5}k'_{3,2}w_{2,2}^2\hat{w}_{2,2}^3 + w_{3,2}\frac{2}{w_{2,2}^4}k'_{2,2}\hat{w}_{2,2}^3 + w_{3,2}w_{2,2}^2\frac{3}{\hat{w}_{2,2}^3}\hat{k}'_{2,2}\Big) \\
&+ y_{13,1}\Big(\frac{4}{w_{3,2}^2}k'_{3,2}w_{2,2}^2 + w_{3,2}^4\frac{2}{w_{2,2}^4}k'_{2,2}\Big) + y_{15,1}\Big(\frac{2}{w_{3,2}^4}k'_{3,2}w_{2,2}^4 + w_{3,2}^2\frac{4}{w_{2,2}^2}k'_{2,2}\Big) \\
&+ y_{14,1}\Big(\frac{3}{w_{3,2}^3}k'_{3,2}\hat{w}_{2,2}^3 + w_{3,2}^3\frac{3}{\hat{w}_{2,2}^3}\hat{k}'_{2,2}\Big) + \Big(\frac{5}{w_{3,2}}k'_{3,2}w_{2,2}^4\hat{w}_{2,2}^3 + w_{3,2}^5\frac{4}{w_{2,2}^2}k'_{2,2}\hat{w}_{2,2}^3 + w_{3,2}^5 w_{2,2}^4\frac{3}{\hat{w}_{2,2}^3}\hat{k}'_{2,2}\Big)\Big) \\
\\
&= y_{16,1}\Big(k'_{3,2}k_{2,2}\hat{k}_{2,2} + 2k_{3,2}k'_{2,2}\hat{k}_{2,2} + 3k_{3,2}k_{2,2}\hat{k}'_{2,2}\Big) \\
&+ y_{13,1}y_{15,2}\Big(4k'_{3,2}k_{2,2} + 2k_{3,2}k'_{2,2}\Big) + y_{15,1}y_{13,2}\Big(2k'_{3,2}k_{2,2} + 4k_{3,2}k'_{2,2}\Big) + y_{14,1}y_{14,2}\Big(3k'_{3,2}\hat{k}_{2,2} + 3k_{3,2}\hat{k}'_{2,2}\Big) \\
&+ y_{16,2}\Big(5k'_{3,2}k_{2,2}\hat{k}_{2,2} + 4k_{3,2}k'_{2,2}\hat{k}_{2,2} + 3k_{3,2}k_{2,2}\hat{k}'_{2,2}\Big).
\end{aligned}
$$

$$
\begin{aligned}
&6y_{13,2}y_{16,2}\frac{\partial A(P_2,P_1)}{\partial x_2}+6y_{13,1}y_{16,1}\frac{\partial A(P_1,P_2)}{\partial x_2}\\
&=y_{16,1}\Big(k'_{3,2}k_{2,2}\hat{k}_{2,2}+2k_{3,2}k'_{2,2}\hat{k}_{2,2}+3k_{3,2}k_{2,2}\hat{k}'_{2,2}+5k'_{3,1}k_{2,1}\hat{k}_{2,1}+4k_{3,1}k'_{2,1}\hat{k}_{2,1}+3k_{3,1}k_{2,1}\hat{k}'_{2,1}\Big)\\
&+y_{13,1}y_{15,2}\Big(4k'_{3,2}k_{2,2}+2k_{3,2}k'_{2,2}+2k'_{3,1}k_{2,1}+4k_{3,1}k'_{2,1}\Big)\\
&+y_{15,1}y_{13,2}\Big(2k'_{3,2}k_{2,2}+4k_{3,2}k'_{2,2}+4k'_{3,1}k_{2,1}+2k_{3,1}k'_{2,1}\Big)\\
&+y_{14,1}y_{14,2}\Big(3k'_{3,2}\hat{k}_{2,2}+3k_{3,2}\hat{k}'_{2,2}+3k'_{3,1}\hat{k}_{2,1}+3k_{3,1}\hat{k}'_{2,1}\Big)\\
&+y_{16,2}\Big(5k'_{3,2}k_{2,2}\hat{k}_{2,2}+4k_{3,2}k'_{2,2}\hat{k}_{2,2}+3k_{3,2}k_{2,2}\hat{k}'_{2,2}+k'_{3,1}k_{2,1}\hat{k}_{2,1}+2k_{3,1}k'_{2,1}\hat{k}_{2,1}+3k_{3,1}k_{2,1}\hat{k}'_{2,1}\Big).
\end{aligned}
$$

Letting the terms of $B(P_1,P_2)$ be denoted by

$$
\begin{aligned}
B(P_1,P_2)&=y_{16,1}B_{16}(P_1,P_2)+y_{13,1}y_{15,2}B_{13,15}(P_1,P_2)+y_{14,1}y_{14,2}B_{14}(P_1,P_2)\\
&\quad+y_{13,2}y_{15,1}B_{13,15}(P_2,P_1)+y_{16,2}B_{16}(P_2,P_1),
\end{aligned}
$$

where B_{16}, $B_{13,14}$ and B_{15} are rational functions of x_1 and x_2, we evaluate:

$$
\begin{aligned}
B_{16}(P_1,P_2)=\frac{(k_{3,2}k_{2,2}\hat{k}_{2,2}-k_{3,1}k_{2,1}\hat{k}_{2,1})}{x_1-x_2}&+k'_{3,2}k_{2,2}\hat{k}_{2,2}+2k_{3,2}k'_{2,2}\hat{k}_{2,2}+3k_{3,2}k_{2,2}\hat{k}'_{2,2}\\
&+5k'_{3,1}k_{2,1}\hat{k}_{2,1}+4k_{3,1}k'_{2,1}\hat{k}_{2,1}+3k_{3,1}k_{2,1}\hat{k}'_{2,1},
\end{aligned}
$$

$$
B_{13,15}(P_1,P_2)=\frac{k_{3,2}k_{2,2}-k_{3,1}k_{2,1}}{x_1-x_2}+2k'_{3,2}k_{2,2}+4k_{3,2}k'_{2,2}-4k'_{3,1}k_{2,1}-2k_{3,1}k'_{2,1},
$$

and

$$
B_{14}(P_1,P_2)=\frac{k_{3,2}\hat{k}_{2,2}-k_{3,1}\hat{k}_{2,1}}{x_1-x_2}+3k'_{3,2}\hat{k}_{2,2}+3k_{3,2}\hat{k}'_{2,2}-3k'_{3,1}\hat{k}_{2,1}-3k_{3,1}\hat{k}'_{2,1}.
$$

We are concerned with $B(P_1,P_2)/(x_1-x_2)$ and thus compute

$$
\begin{aligned}
\frac{B_{16}(P_1,P_2)}{(x_1-x_2)}&=-7x_2^5+\Big(-6\lambda_1^{(3)}-x_1-5\lambda_1^{(2)}-4\hat{\lambda}_1^{(2)}\Big)x_2^4\\
&+\Big(5x_1^2+(2\hat{\lambda}_1^{(2)}+\lambda_1^{(2)})x_1-3\lambda_2^{(2)}-3\lambda_1^{(3)}\hat{\lambda}_1^{(2)}-\hat{\lambda}_2^{(2)}-5\lambda_2^{(3)}-2\lambda_1^{(2)}\hat{\lambda}_1^{(2)}-4\lambda_1^{(3)}\lambda_1^{(2)}\Big)x_2^3\\
&+\Big(11x_1^3+(6\lambda_1^{(3)}+7\lambda_1^{(2)}+8\hat{\lambda}_1^{(2)})x_1^2+(4\lambda_1^{(2)}\hat{\lambda}_1^{(2)}+3\lambda_1^{(3)}\hat{\lambda}_1^{(2)}+2\lambda_1^{(3)}\lambda_1^{(2)}\\
&\quad+\lambda_2^{(3)}+3\lambda_2^{(2)}+5\hat{\lambda}_2^{(2)})x_1-\lambda_1^{(3)}\lambda_1^{(2)}\hat{\lambda}_1^{(2)}-2\lambda_2^{(3)}\hat{\lambda}_1^{(2)}-3\lambda_2^{(3)}\lambda_1^{(2)}-2\lambda_1^{(3)}\lambda_2^{(2)}\\
&\quad+\lambda_1^{(2)}\hat{\lambda}_2^{(2)}-4\lambda_3^{(3)}\Big)x_2^2
\end{aligned}
$$

$$
\begin{aligned}
+\Big(&17x_1^4 + (13\lambda_1^{(2)} + 12\lambda_1^{(3)} + 14\hat{\lambda}_1^{(2)})x_1^3 \\
&+ (10\lambda_1^{(2)}\hat{\lambda}_1^{(2)} + 9\lambda_1^{(3)}\hat{\lambda}_1^{(2)} + 8\lambda_1^{(3)}\lambda_1^{(2)} + 11\hat{\lambda}_2^{(2)} + 9\lambda_2^{(2)} + 7\lambda_2^{(3)})x_1^2 \\
&+ (7\lambda_1^{(2)}\hat{\lambda}_2^{(2)} + 6\lambda_1^{(3)}\hat{\lambda}_2^{(2)} + 4\lambda_1^{(3)}\lambda_2^{(2)} + 6\lambda_2^{(2)}\hat{\lambda}_1^{(2)} + 4\lambda_2^{(3)}\hat{\lambda}_1^{(2)} + 3\lambda_2^{(3)}\lambda_1^{(2)} \\
&\quad + 2\lambda_3^{(3)} + 5\lambda_1^{(3)}\lambda_1^{(2)}\hat{\lambda}_1^{(2)})x_1 \\
&+ \lambda_1^{(3)}\lambda_2^{(2)}\hat{\lambda}_1^{(2)} + 2\lambda_1^{(3)}\lambda_1^{(2)}\hat{\lambda}_2^{(2)} + 3\lambda_2^{(2)}\hat{\lambda}_2^{(2)} - 2\lambda_3^{(3)}\lambda_1^{(2)} - \lambda_2^{(3)}\lambda_2^{(2)} \\
&+ \lambda_2^{(3)}\hat{\lambda}_2^{(2)} - \lambda_3^{(3)}\hat{\lambda}_1^{(2)}\Big)x_2
\end{aligned}
$$

$$
\begin{aligned}
&+ 23x_1^5 + (19\lambda_1^{(2)} + 18\lambda_1^{(3)} + 20\hat{\lambda}_1^{(2)})x_1^4 + (14\lambda_1^{(3)}\lambda_1^{(2)} + 15\lambda_1^{(3)}\hat{\lambda}_1^{(2)} + 16\lambda_1^{(2)}\hat{\lambda}_1^{(2)} \\
&\quad + 13\lambda_2^{(3)} + 15\lambda_2^{(2)} + 17\hat{\lambda}_2^{(2)})x_1^3 \\
&\quad + (10\lambda_1^{(3)}\lambda_2^{(2)} + 9\lambda_2^{(3)}\lambda_1^{(2)} + 10\lambda_2^{(3)}\hat{\lambda}_1^{(2)} + 12\lambda_2^{(2)}\hat{\lambda}_1^{(2)} + 12\lambda_1^{(3)}\hat{\lambda}_2^{(2)} + 13\lambda_1^{(2)}\hat{\lambda}_2^{(2)} \\
&\quad + 8\lambda_3^{(3)} + 11\lambda_1^{(3)}\lambda_1^{(2)}\hat{\lambda}_1^{(2)})x_1^2 \\
&\quad + (4\lambda_3^{(3)}\lambda_1^{(2)} + 5\lambda_2^{(3)}\lambda_2^{(2)} + 5\lambda_3^{(3)}\hat{\lambda}_1^{(2)} + 7\lambda_2^{(3)}\hat{\lambda}_2^{(2)} + 9\lambda_2^{(2)}\hat{\lambda}_2^{(2)} + 6\lambda_2^{(3)}\lambda_1^{(2)}\hat{\lambda}_1^{(2)} \\
&\quad + 8\lambda_1^{(3)}\lambda_1^{(2)}\hat{\lambda}_2^{(2)} + 7\lambda_1^{(3)}\lambda_2^{(2)}\hat{\lambda}_1^{(2)})x_1 \\
&\quad + 3\lambda_2^{(3)}\lambda_1^{(2)}\hat{\lambda}_2^{(2)} + 4\lambda_1^{(3)}\lambda_2^{(2)}\hat{\lambda}_2^{(2)} + 2\lambda_2^{(3)}\lambda_2^{(2)}\hat{\lambda}_1^{(2)} + \lambda_3^{(3)}\lambda_1^{(2)}\hat{\lambda}_1^{(2)} + 2\lambda_3^{(3)}\hat{\lambda}_2^{(2)}.
\end{aligned}
$$

$$
\begin{aligned}
\frac{B_{13,15}(P_1,P_2)}{(x_1-x_2)} &= -8x_2^3 + (-4\lambda_1^{(2)} - 6\lambda_1^{(3)} - 2x_1)x_2^2 + (-2\lambda_1^{(3)}\lambda_1^{(2)} + 2\lambda_1^{(2)}x_1 - 4\lambda_2^{(3)} + 4x_1^2)x_2 \\
&\quad + 10x_1^3 + (8\lambda_1^{(2)} + 6\lambda_1^{(3)})x_1^2 + (4\lambda_1^{(3)}\lambda_1^{(2)} + 2\lambda_2^{(3)} + 6\lambda_2^{(2)})x_1 + 2\lambda_1^{(3)}\lambda_2^{(2)} - 2\lambda_3^{(3)}, \\
\frac{B_{14}(P_1,P_2)}{(x_1-x_2)} &= -9x_2^3 + (-6\hat{\lambda}_1^{(2)} - 3x_1 - 6\lambda_1^{(3)})x_2^2 + (3x_1^2 - 3\lambda_1^{(3)}\hat{\lambda}_1^{(2)} \\
&- 3\hat{\lambda}_2^{(2)} - 3\lambda_2^{(3)})x_2 + 9x_1^3 + (6\lambda_1^{(3)} + 6\hat{\lambda}_1^{(2)})x_1^2 + (3\lambda_2^{(3)} + 3\lambda_1^{(3)}\hat{\lambda}_1^{(2)} + 3\hat{\lambda}_2^{(2)})x_1.
\end{aligned}
$$

By these results we obtain the $\nu^{II;12}$'s in the statement. □

Corollary 4.7. *(1) The one-form*

$$\Pi^{(12)}{}_{P_1}^{P_2}(P) := \Sigma^{(12)}(P, P_1)dx - \Sigma^{(12)}(P, P_2)dx$$

is a differential of the third kind, whose only (first-order) poles are $P = P_1$ and $P = P_2$, and residues $+1$ and -1 respectively.

(2) The fundamental differential of the second kind $\Omega^{(12)}(P_1, P_2)$ is given by

$$\begin{aligned}\Omega^{(12)}(P_1, P_2) &= d_{P_2}\Sigma^{(12)}(P_1, P_2) + \sum_{i=1}^{12} \nu_i^{I;12}(P_1) \otimes \nu_i^{II;12}(P_2) \\ &= \frac{F^{(12)}(P_1, P_2) dx_1 \otimes dx_2}{(x_1 - x_2)^2 6^2 y_{13,P_1} y_{16,P_1} y_{13,P_2} y_{16,P_2}},\end{aligned} \tag{4.17}$$

where $F^{(4)}$ is an element of $R_{12} \otimes R_{12}$.

Lemma 4.8. *We have*

$$\lim_{P_1 \to \infty} \frac{F^{(12)}(P_1, P_2)}{\phi_{11}^{(12)}(P_1)(x_1 - x_2)^2} = \phi_{12}^{(12)}(P_2) = x_{P_2}^4. \tag{4.18}$$

For later convenience we introduce the notation:

$$\begin{aligned}\Omega^{(12)}{}_{Q_1,Q_2}^{P_1,P_2} &:= \int_{P_2}^{P_1} \int_{Q_2}^{Q_1} \Omega^{(12)}(P, Q) \\ &= \int_{P_2}^{P_1} (\Sigma^{(12)}(P, Q_1) - \Sigma^{(12)}(P, Q_2)) + \sum_{i=1}^{12} \int_{P_2}^{P_1} \nu_i^{I;12}(P) \int_{Q_2}^{Q_1} \nu_i^{II;12}(P).\end{aligned} \tag{4.19}$$

5. The sigma function over $(3, 7, 8)$ and $(6, 13, 14, 15, 16)$ curves

5.1. **Legendre relation.** In this section, g will be 4 or 12. In the monomial notation, as there were two conventions for X_4 and X_{12}, $\phi^{(g)}$ will be $\phi_{H^1}^{(4)}$ for $g = 4$ and $\phi^{(12)}$ for $g = 12$.

We write the periods:

$$\left[\eta^{(g)\prime} \ \eta^{(g)\prime\prime}\right] := \frac{1}{2}\left[\int_{\alpha_i^{(g)}} \nu_j^{II;g} \quad \int_{\beta_i^{(g)}} \nu_j^{II;g}\right]_{i,j=1,2,\cdots,g}. \tag{5.1}$$

Let $\tau_{Q_1,Q_2}^{(g)}$ be the normalized differential of the third kind that has residues $+1$, -1 at Q_1, Q_2, is regular everywhere else, and is normalized, $\int_{\alpha_i} \tau_{P,Q}^{(g)} = 0$ for every i [FK, III.3.5]. The following Lemma corresponding to Corollary 2.6 (ii) in [Fay] holds:

Lemma 5.1.

$$\Omega^{(g)}{}_{Q_1,Q_2}^{P_1,P_2} = \int_{P_2}^{P_1} \tau_{Q_1,Q_2}^{(g)} + \sum_{i,j=1}^{g} \gamma_{ij}^{(g)} \int_{P_2}^{P_1} \nu_i^{I;g} \int_{Q_2}^{Q_1} \nu_j^{I;g},$$

where $\gamma^{(g)} = \omega^{(g)\prime -1} \eta^{(g)\prime}$.

Proof. $\partial_{P_1}\Omega^{(g)}{}^{P_1,P_2}_{Q_1,Q_2} - \tau^{(g)}_{Q_1,Q_2}$ must be expressed by the linear combination of the holomorphic one forms. Noting $\int_{P_1}^{P_2}\tau^{(g)}_{Q_1,Q_2} = \int_{Q_1}^{Q_2}\tau^{(g)}_{P_1,P_2}$ [FK, III.3.5], $\gamma^{(g)}_{ij}$ is symmetric. Due to (3.16) and (4.19),

$$(\Sigma(P_1,Q_1)-\Sigma(P_1,Q_2))+\sum_{i=1}^{g}\nu_i^{I;g}(P_1)\int_{Q_2}^{Q_1}\nu_i^{II;g}(P) = \tau^{(g)}_{Q_1,Q_2}(P_1)+\sum_{i,j=1}^{g}\gamma^{(g)}_{ij}\nu_i^{I;g}(P_1)\int_{Q_2}^{Q_1}\nu_j^{I;g}.$$

By choosing an appropriate path Γ from Q_1 to Q_2, homotopic to $\alpha^{(g)}_k$, we have $(\eta^{(g)\prime}\nu^{I;g})^t = \gamma^{(g)}\cdot(\omega^{(g)\prime}\nu^{I;g})^t$. □

The following Proposition provides *generalized Legendre relation* [B1, BLE1, BLE2], which determines a symplectic structure in the Jacobian [Po, Chap.3,4].

Proposition 5.2. (generalized Legendre relation) *The matrix,*

$$M := \begin{bmatrix} 2\omega^{(g)\prime} & 2\omega^{(g)\prime\prime} \\ 2\eta^{(g)\prime} & 2\eta^{(g)\prime\prime} \end{bmatrix}, \tag{5.2}$$

satisfies

$$M\begin{bmatrix} & -1 \\ 1 & \end{bmatrix}{}^tM = 2\pi\sqrt{-1}\begin{bmatrix} & -1 \\ 1 & \end{bmatrix}. \tag{5.3}$$

Proof. By comparing Lemma 5.1 and, (3.14) in Corollary 3.12 and (4.17) in Corollary 4.7, we choose appropriately $(2g)^2$ paths and take the integrals along these paths. For example, for

$$\int_{P_2}^{P_1}(\Sigma(P,Q_1)-\Sigma(P,Q_2))+\sum_{i=1}^{g}\int_{P_2}^{P_1}\nu_i^{I;g}(P)\int_{Q_2}^{Q_1}\nu_i^{II;g}(P)$$

$$=\int_{Q_2}^{Q_1}\tau^{(g)}_{P_1,P_2}+\sum_{i,j=1}^{g}\gamma^{(g)}_{ij}\int_{P_2}^{P_1}\nu_i^{I;g}\int_{Q_2}^{Q_1}\nu_j^{I;g},$$

the contour of Q_1 to Q_2 along α_k gives

$$\sum_{i=1}^{g}\int_{P_2}^{P_1}\nu_i^{I;g}(P)\eta^{(g)\prime}_{ki}(P) = \sum_{i,j=1}^{g}\gamma^{(g)}_{ij}\int_{P_2}^{P_1}\nu_i^{I;g}\omega^{(g)\prime}_{kj},$$

and further contour integration provides ${}^t({}^t\omega^{(g)\prime}\eta^{(g)\prime}) = ({}^t\omega^{(g)\prime}\eta^{(g)\prime})$. The contour of Q_1 to Q_2 along $\beta^{(g)}_k$ gives

$$\sum_{i=1}^{g}\int_{P_2}^{P_1}\nu_i^{I;g}(P)\eta^{(g)\prime\prime}_{ki}(P) = 2\pi\sqrt{-1}\int_{P_2}^{P_1}({\omega^{(g)\prime}}^{-1}\nu^{I;g})_k+\sum_{i,j=1}^{g}\gamma^{(g)}_{ij}\int_{P_2}^{P_1}\nu_i^{I;g}\omega^{(g)\prime}_{kj},$$

which, in view of the contour integral, turns out to be

$${}^t({}^t\omega^{(g)\prime\prime}\eta^{(g)\prime\prime}) = 2\pi\sqrt{-1}\mathbb{T}+({}^t\omega^{(g)\prime\prime}\gamma^{(g)}\omega^{(g)\prime\prime}),$$

$$({}^t\omega^{(g)\prime}\eta^{(g)\prime\prime}) = 2\pi\sqrt{-1}1_g + ({}^t\omega^{(g)\prime}\gamma^{(g)}\omega^{(g)\prime\prime}).$$

Here we use $\int_{\beta_k} \tau^{(g)}_{Q_1,Q_2} = 2\pi\sqrt{-1}\int_{Q_2}^{Q_1}(\omega^{(g)\prime-1}\nu^{I;g})_k$ [FK, III.3.5]. □

5.2. σ **function.** By the Riemann relations [Fay], it is known that $\mathrm{Im}\,(\omega^{(g)\prime^{-1}}\omega^{(g)\prime\prime})$ is positive definite. As in Theorem 1.1 in [Fay], let

$$\delta := \left[\begin{array}{c} \delta' \\ \delta'' \end{array}\right] \in \left(\tfrac{1}{2}\mathbb{Z}\right)^{2g} \tag{5.4}$$

be the theta characteristic which is equal to the Riemann constant ξ_R with respect to the base point ∞ and the period matrix $[\,2\omega^{(g)\prime}\ 2\omega^{(g)\prime\prime}]$.

We define an entire function of (a column-vector) $u = {}^t(u_1, u_2, \dots, u_g) \in \mathbb{C}^g$,

(5.5)

$$\begin{aligned}
\sigma^{(g)}(u) &= \sigma^{(g)}(u; M) = \sigma^{(g)}(u_1, u_2, \dots, u_g; M)\\
&= c\exp(-\tfrac{1}{2}\ {}^t u\eta^{(g)\prime}\omega^{(g)\prime^{-1}}u)\vartheta[\delta]\,(\frac{1}{2}\omega^{(g)\prime^{-1}}u;\ \omega^{(g)\prime^{-1}}\omega^{(g)\prime\prime})\\
&= c\exp(-\tfrac{1}{2}\ {}^t u\eta^{(g)\prime}\omega^{(g)\prime^{-1}}\ u)\\
&\qquad \times \sum_{n\in\mathbb{Z}^g} \exp\left[\pi\sqrt{-1}\{\ {}^t(n+\delta')\omega^{(g)\prime^{-1}}\omega^{(g)\prime\prime}(n+\delta') +\ {}^t(n+\delta')(\omega^{(g)\prime^{-1}}u+\delta'')\}\right],
\end{aligned}$$

where c is a certain constant of a rational of b's.

For a given $u \in \mathbb{C}^g$, we introduce u' and u'' in $\mathbb{R}^g$ so that

$$u = 2\omega^{(g)\prime}u' + 2\omega^{(g)\prime\prime}u''.$$

Proposition 5.3. *For* $u,\ v \in \mathbb{C}^g$*, and* $\ell\ (= 2\omega^{(g)\prime}\ell' + 2\omega^{(g)\prime\prime}\ell'') \in \Lambda$*, we define*

$$\begin{aligned}
L(u, v) &:= 2\ {}^t u(\eta' v' + \eta'' v''),\\
\chi(\ell) &:= \exp[\pi\sqrt{-1}\big(2({}^t\ell'\delta'' - {}^t\ell''\delta') + {}^t\ell'\ell''\big)]\ (\in \{1,\ -1\}).
\end{aligned}$$

The following holds

$$\sigma^{(g)}(u+\ell) = \sigma^{(g)}(u)\exp(L(u+\frac{1}{2}\ell, \ell))\chi(\ell). \tag{5.6}$$

Proof. The proof is standard, noting that $\sigma^{(g)}$ is associated to the normalized theta function in Chapter VI of [L]. □

Remark 5.4. Following a formula proven for the rational/polynomial case by Buchstaber, Leykin and Enolskii [BEL2], in [Na1] Nakayashiki showed that the leading of the σ function associated with (r, s) curve is expressed by a Schur function by normalizing the constant factor c. We also expect that

$$\sigma^{(g)}(u) = S_\Lambda(T)|_{T_{\Lambda_i+g-i}=u_i} + \sum_\alpha a_\alpha u^\alpha$$

where $a_\alpha \in \mathbb{Q}[b_1, \cdots, b_\ell]$ for $\ell = 5$ for X_4 and $\ell = 7$ for X_{12}, $\alpha = \alpha_1 \cdots \alpha_g$ and $u^\alpha = u_1^{\alpha_1} \cdots u_g^{\alpha_g}$. Here for a Young diagram Λ, S_Λ and s_Λ are the Schur functions defined by

$$S_\Lambda(T) = s_\Lambda(t), \quad T_k := \frac{1}{k} \sum_{i=1}^{g} t_j^k.$$

The vanishing locus of $\sigma^{(g)}$ is:

$$\Theta^{g-1} = (\mathcal{W}^{g-1} \cup [-1]\mathcal{W}^{g-1}) = \mathcal{W}^{g-1}. \tag{5.7}$$

The last equality is due to Proposition 5.3, which shows that $\sigma^{(g)}$ is an even or odd function under the action of $[-1]$; the reason for introducing $\mathcal{W}^{g-1} \cup [-1]\mathcal{W}^{g-1}$ is that the analogous loci when $g-1$ is replaced by k play an important role and $\mathcal{W}^k$ is not $[-1]$-invariant in general.

5.3. **The Riemann fundamental relation** [R]**.** We review a relation which we call the Riemann fundamental relation [R], [B1, §195]:

Proposition 5.5. *For* $(P, Q, P_i, P'_i) \in X^2 \times (S^g(X) \setminus S^g_1(X)) \times (S^g(X) \setminus S^g_1(X))$,

$$u := \sum_{i=1}^{g} \hat{u}(P_i), \quad v := \sum_{i=1}^{g} \hat{u}(P'_i),$$

$$\begin{aligned} \exp\left(\sum_{i,j=1}^{g} \Omega^{(g)}{}^{P,Q}_{P_i,P'_j} \right) &= \frac{\sigma^{(g)}(\hat{u}(P) - u)\sigma^{(g)}(\hat{u}(Q) - v)}{\sigma^{(g)}(\hat{u}(Q) - u)\sigma^{(g)}(\hat{u}(P) - v)} \\ &= \frac{\sigma^{(g)}(\hat{u}(P) - \hat{u}(P_1, \cdots, P_g))\sigma^{(g)}(\hat{u}(Q) - \hat{u}(P'_1, \cdots, P'_g))}{\sigma^{(g)}((\hat{u}(Q) - \hat{u}(P_1, \cdots, P_g))\sigma^{(g)}(\hat{u}(P) - \hat{u}(P'_1, \cdots, P'_g))}. \end{aligned}$$

Proof. The right-hand side can be expressed as

$$\exp(\text{bilinear term in } w\text{'s}) \frac{\theta(\omega^{(g)\prime -1}(\hat{u}(P) - u) + \xi_R)\theta(\omega^{(g)\prime -1}(\hat{u}(Q) - v) + \xi_R)}{\theta(\omega^{(g)\prime -1}(\hat{u}(Q) - u) + \xi_R)\theta(\omega^{(g)\prime -1}(\hat{u}(P) - v)) + \xi_R)},$$

where ξ_R is the Riemann constant. By Riemann's theorem for theta functions [Fay, p.23], the above becomes

$$\exp(\text{bilinear term in } w\text{'s}) \exp\left(\sum_{j=1}^{g} \int_Q^P \tau^{(g)}_{P_j, P'_j} \right).$$

The exponential part of the bilinear term is given by,

$$(u - v)^t \gamma^{(g)}(\hat{u}(P) - \hat{u}(Q)) = \sum_{i,j,k} \gamma^{(g)}_{ij} \int_{P'_k}^{P_k} \nu_i^{I;g} \int_Q^P \nu_j^{I;g},$$

which equals (Lemma (5.1))

$$\sum_{i=1}^{g} \Omega^{(g)}{}^{P,Q}_{P_i,P_i'} - \sum_{i=1}^{g} \int_Q^P \tau^{(g)}_{P_i,P_i'}.$$

The above integrals depend upon the paths we choose, but (5.6) shows that such dependence cancels. Thus the right-hand side coincides with the left-hand side. □

Proposition 5.6. *For* $(P, P_1, \ldots, P_g) \in X \times S^g(X) \setminus S^g_1(X)$ *and* $u := \hat{u}(P_1, \ldots, P_g)$*, the equality*

$$\sum_{i,j=1}^{g} \wp^{(g)}_{i,j}\left(\hat{u}(P) - u\right) \phi^{(g)}_{i-1}(P) \phi^{(g)}_{j-1}(P_a) = \frac{F^{(g)}(P, P_a)}{(x - x_a)^2},$$

holds for every $a = 1, 2, \ldots, g$*, where we set*

$$\wp^{(g)}_{ij}(u) := -\frac{\sigma^{(g)}_i(u)\sigma^{(g)}_j(u) - \sigma^{(g)}(u)\sigma^{(g)}_{ij}(u)}{\sigma^{(g)}(u)^2} \equiv -\frac{\partial^2}{\partial u_i \partial u_j} \log \sigma^{(g)}(u).$$

Here for $g = 4$ *case,* $\phi^{(g)}_i$ *is interpreted as* $\phi^{(4)}_{H^1 i}$*.*

Proof. For the case X_4, using the relation (3.8) and taking logarithm of both sides in the Riemann fundamental relation and differentiating along $P_1 = P$ and $P_2 = P_a$, we obtain the claim. Similarly we have the result for the case X_{12} using (4.11) □

6. JACOBI INVERSION FORMULAE FOR $(3, 7, 8)$ AND $(6, 13, 14, 15, 16)$ CURVES

In this section, we will consider a Jacobi-inversion type formulae associated with X_4 and X_{12}.

6.1. **The $\mu^{(g)}_n$ functions over $(3, 7, 8)$ and $(6, 13, 14, 15, 16)$ curves.** In this subsection, g will be 4 or 12, and $\phi^{(g)}$ accordingly, $\phi^{(4)}_{H^1}$, $\phi^{(4)}_{H^0}$, or $\phi^{(12)}$. In [MP08], we introduced meromorphic functions on the curve, reviewed here in Definition 6.1, which generalize the polynomial U in Mumford's (U, V, W) parameterization of a hyperelliptic Jacobian (which he attributes to Jacobi) [Mum].

For the definition of the function μ's, we introduce the Frobenius-Stickelberger (FS) matrix and its determinant following [MP08]. Let n be a positive integer and $P_1, \ldots, P_n$ be in $X_g \backslash \infty$. As in (3.7) and (4.10), we define the *ℓ-reduced Frobenius-Stickelberger* (FS) *matrix* by:

$$\Psi^{(g;\check{\ell})}_n(P_1, P_2, \ldots, P_n) := \begin{pmatrix} \phi^{(g)}_0(P_1) & \phi^{(g)}_1(P_1) & \phi^{(g)}_2(P_1) & \cdots & \check{\phi^{(g)}_\ell}(P_1) & \cdots & \phi^{(g)}_n(P_1) \\ \phi^{(g)}_0(P_2) & \phi^{(g)}_1(P_2) & \phi^{(g)}_2(P_2) & \cdots & \check{\phi^{(g)}_\ell}(P_2) & \cdots & \phi^{(g)}_n(P_2) \\ \vdots & \vdots & \vdots & \ddots & \vdots & \ddots & \vdots \\ \phi^{(g)}_0(P_n) & \phi^{(g)}_1(P_n) & \phi^{(g)}_2(P_n) & \cdots & \check{\phi^{(g)}_\ell}(P_n) & \cdots & \phi^{(g)}_n(P_n) \end{pmatrix}.$$

and $\psi_n^{(g;\check{\ell})}(P_1, P_2, \dots, P_n) := \det \Psi_n^{(g;\check{\ell})}(P_1, P_2, \dots, P_n)$ (a check on top of a letter signifies deletion). It is also convenient to introduce the simpler notation:

$$\psi_n^{(g)}(P_1, \dots, P_n) := \det(\Psi_n^{(g;\check{n})}(P_1, \dots, P_n)), \quad \Psi_n^{(g)}(P_1, \dots, P_n) := \Psi_n^{(g;\check{n})}(P_1, \dots, P_n),$$

for the un-bordered matrix. We call this matrix *Frobenius-Stickelberger (FS) matrix* and its determinant *Frobenius-Stickelberger (FS) determinant.* These become singular for some tuples in $(X_g \backslash \infty)^n$.

Definition 6.1. *For* $P, P_1, \dots, P_n \in (X_g \backslash \infty) \times SS^n(X_g \backslash \infty)$*, we define* $\mu_n^{(g)}(P)$ *by*

$$\mu_n^{(g)}(P) := \mu_n^{(g)}(P; P_1, \dots, P_n) := \lim_{P'_i \to P_i} \frac{1}{\psi_n^{(g)}(P'_1, \dots, P'_n)} \psi_{n+1}^{(g)}(P'_1, \dots, P'_n, P),$$

where the P'_i *are generic, the limit is taken (irrespective of the order) for each* i*; and* $\mu_{n,k}^{(g)}(P_1, \dots, P_n)$ *by*

$$\mu_n^{(g)}(P) = \phi_n^{(g)}(P) + \sum_{k=0}^{n-1} (-1)^{n-k} \mu_{n,k}^{(g)}(P_1, \dots, P_n) \phi_k^{(g)}(P),$$

with the convention $\mu_{n,n}^{(g)}(P_1, \dots, P_n) \equiv 1$*. Let* $\mu_{H^a}^{(4)}$ *be* $\mu^{(4)}$ *for* $\phi_{H^a}^{(4)}$ *for* $a = 0, 1$*.*

Meromorphic functions, viewed as divisors on the curve, allow us to express the addition structure of $\mathrm{Pic} X_g$ in terms of FS-matrices. For n points $(P_i)_{i=1,\dots,n} \in X_g \backslash \infty$, we find an element of R associated with any point $P = (x, y)$ in $(X_g \backslash \infty)$, $\alpha_n(P) := \alpha_n(P; P_1, \dots, P_n) = \sum_{i=0}^n a_i \phi_i(P)$, $a_i \in \mathbb{C}$ and $a_n = 1$, which has a zero at each point P_i (with multiplicity, if the P_i are repeated) and has smallest possible order of pole at ∞ with this property. α_n can be identified with $\mu_{H^0,n}^{(4)}$ for X_4 and $\mu_n^{(12)}$ for X_{12}.

On X_4, instead of the pole condition on elements of $H^0(X_4 \setminus \infty, \mathcal{O}_{X_4})$, we impose an alternative condition to determine a polynomial $\alpha_{H^1 n} = \sum_{i=0}^n a_i \phi_{H^1 i}^{(4)}(P)$, $a_i \in \mathbb{C}$ and $a_n = 1$, namely to have a (possibly multiple) zero at each point P_i and smallest possible degree to match the order of pole of $dx/3y_7y_8$ at ∞; $\alpha_{H^1 n}$ can be identified with $\mu_{H^1,n}^{(4)}$, in fact $\dfrac{\mu_{H^1,n}^{(4)} dx}{3y_7y_8}$ does not have a singularity over $X \setminus \infty$ whereas $\dfrac{\mu_{H^0,n}^{(4)} dx}{3y_7y_8}$ does.

For α_n for X_g ($g = 4, 12$) and $\alpha_{H^1 n}$ for X_4 and $N^{(g)}(n)$ $(N_{H^1}^{(4)}(n), N_{H^0}^{(4)}(n), N^{(12)}(n))$, we have the following lemma.

Lemma 6.2. *Let* n *be a positive integer. For* $(P_i)_{i=1,\dots,n} \in SS^n(X \backslash \infty)$*, the function* α_n *over* X_g *induces the map (which we call by the same name):*

$$\alpha_n : SS^n(X_g \backslash \infty) \to SS^{N^{(g)}(n)-n}(X_g),$$

i.e., to $(P_i)_{i=1,\dots,n} \in SS^n(X_g \backslash \infty)$ *there corresponds an element* $(Q_i)_{i=1,\dots,N^{(g)}(n)-n} \in SS^{N^{(g)}(n)-n}(X_g)$, *such that*

$$\sum_{i=1}^{n} P_i - n\infty \sim - \sum_{i=1}^{N^{(g)}(n)-n} Q_i + (N^{(g)}(n) - n)\infty.$$

We want the preimage of α_n to include the base point ∞. For an effective divisor D in $SS^n(X_g)$ of degree n, let D' be the maximal subdivisor of D which does not contain ∞, $D = D' + (n-m)\infty$ where $\deg D' = m(\leq n)$ and $D' \in SS^m(X_g \backslash \infty)$. Then we extend the map to $\overline{\alpha}_n$ by defining $\overline{\alpha}_n(D) = \alpha_m(D') + [N^{(g)}(n) - n - (N^{(g)}(m) - m)]\infty$. In [MP09], we use the following relation as the Serre duality but due to Remark 3.5, the expression should be modified for X_4.

We see from the linear equivalence of Lemma 6.2:

Proposition 6.3. *For a positive integer, the Abel map composed with* α_n ($\mu^{(12)}$ *for* X_{12} *and* $\mu^{(4)}_{H^0}$, *for* X_4), *and* $\alpha_{H^1 n}$ *for* $\mu^{(4)}_{H^1}$, *induce*

$$\iota_n : \mathcal{W}^n \to \mathcal{W}^{N^{(g)}(n)-n},$$

and

$$\iota_{H^1 n} : \mathcal{W}^n \to \mathcal{W}^{N^{(4)}_{H^1}(n)-n}.$$

Let image(ι_n) be denoted by $[-1]\mathcal{W}^n$, especially $\iota_g : \mathcal{W}^g \to [-1]\mathcal{W}^g$. Noting for $n \geq g$, $\iota_g \circ \iota_n$ gives the Abel sum

$$\mathcal{W}^n \xrightarrow{\iota_n} \mathcal{W}^g \xrightarrow{\iota_g} \mathcal{W}^g, \quad (\hat{u}(P_1, \dots, P_n) \equiv \hat{u}(Q_1, \dots, Q_g)) \bmod \Lambda.$$

In particular, the addition law on the Jacobian is given by $\iota_g \circ \iota_{2g}$

$$\mathcal{W}^{2g} \xrightarrow{\iota_{2g}} \mathcal{W}^g \xrightarrow{\iota_g} \mathcal{W}^g, \quad (\hat{u}(P_1, \dots, P_g, P'_1, \dots, P'_g) \equiv \hat{u}(Q_1, \dots, Q_g)) \bmod \Lambda.$$

For X_4, every n, $N^{(4)}_{H^1}(n) - n > g \equiv 4$ and thus,

$$\iota_4 \circ \iota_{N^{(4)}_{H^1}(n)} \circ \iota_{H^1 n} : \mathcal{W}^n \to \mathcal{W}^4,$$

which also gives $[-1]$ of $\mathcal{W}_n \to [-1]\mathcal{W}_n \subset \mathcal{W}_4$.

Similar relations hold for $\mu^{(4)}_{H^1}$. We now state the main theorem on the σ-function in the present work.

6.2. Jacobi inversion formulae over Θ^k.

Theorem 6.4. (Jacobi inversion formula)

(1) For $(P, P_1, \cdots, P_4) \in X_4 \times S^4(X_4) \setminus S^4_1(X_4)$, *we have*
  (a) $\mu^{(4)}_{H^1,4}(P; P_1, \dots, P_4) = \phi^{(4)}_{H^1 4}(P) - \sum_{j=1}^{4} \wp^{(4)}_{4j}(\hat{u}(P_1, \dots, P_4))\phi^{(4)}_{H^1 j-1}(P)$.
  (b) $\wp^{(4)}_{4,k+1}(\hat{u}(P_1, \dots, P_4)) = (-1)^{3-k}\mu^{(4)}_{H^1,4,k}(P_1, \dots, P_4), \quad (k = 0, \dots, 3)$.

(2) For $(P, P_1, \cdots, P_{12}) \in X_{12} \times S^{12}(X_{12}) \setminus S^{12}_1(X_{12})$, *we have*

(a) $\mu^{(12)}_{12}(P;P_1,\dots,P_{12}) = \phi^{(12)}_{12}(P) - \sum_{j=1}^{12} \wp^{(12)}_{12,j}(\hat{u}(P_1,\dots,P_{12}))\phi^{(12)}_{j-1}(P)$.

(b) $\wp^{(12)}_{12,k+1}(\hat{u}(P_1,\dots,P_{12})) = (-1)^{12-k-1}\mu^{(12)}_{12,k}(P_1,\dots,P_{12}), \quad (k = 0,\dots,12)$.

Proof. Same as in [MP08, Prop. 4.6]. □

We introduce

$$\Theta^k := \mathcal{W}^k \cup [-1]\mathcal{W}^k. \tag{6.1}$$

Theorem 6.5. *The following relations for $\mu^{(4)}_{H^1,}$ to X_4, and for $\mu^{(12)}$ to X_{12} hold.*

(1) Θ^g *case: for* $(P_1,\dots,P_g) \in S^g(X_g) \setminus S^g_1(X_g)$ *and* $u = \hat{u}(P_1,\dots,P_g) \in \kappa^{-1}(\Theta^g)$,

$$\frac{\sigma^{(g)}_i(u)\sigma^{(g)}_g(u) - \sigma^{(g)}_{gi}(u)\sigma^{(g)}(u)}{{\sigma^{(g)}}^2(u)} = (-1)^{g+1-i}\mu^{(g)}_{g,i-1}(P_1,\dots,P_g), \quad \textit{for } 0 < i \le g.$$

(2) Θ^{g-1} *case: for* $(P_1,\dots,P_{g-1}) \in S^{g-1}(X_g) \setminus S^{g-1}_1(X_g)$ *and* $u = \pm\hat{u}(P_1,\dots,P_{g-1}) \in \kappa^{-1}(\Theta^{g-1})$,

$$\frac{\sigma^{(g)}_i(u)}{\sigma^{(g)}_g(u)} = \begin{cases} (-1)^{g-i}\mu^{(g)}_{g-1,i-1}(P_1,\dots,P_{g-1}) & \textit{for } 0 < i < g, \\ 1 & \textit{for } i = g. \end{cases}$$

(3) Θ^k *case: for* $(P_1,\dots,P_k) \in S^k(X_g) \setminus S^k_1(X_g)$ *and* $u = \pm\hat{u}(P_1,\dots,P_k) \in \kappa^{-1}(\Theta^k)$,

$$\frac{\sigma^{(g)}_i(u)}{\sigma^{(g)}_{k+1}(u)} = \begin{cases} (-1)^{k-i+1}\mu^{(g)}_{k,i-1}(P_1,\dots,P_k) & \textit{for } 0 < i \le k, \\ 1 & \textit{for } i = k+1, \\ 0 & \textit{for } k+1 < i \le g. \end{cases}$$

Proof. Essentially the same as in [MP08, Th. 5.1]. □

We should note that for $k = 1$ case, we have

$$\frac{\sigma^{(12)}_1(u)}{\sigma^{(12)}_2(u)} = -x, \quad \text{for} \quad (x, y_{13}, y_{14}, y_{15}, y_{16}) \in X_{12}.$$

7. Configurations of Curves and Monstrous Moonshine

We highlight some relationships between the curves X_4 and X_{12}, and an additional smooth curve $\hat{X}_4$ of genus 4, with affine plane model of type $(3,4)$, and their algebraic functions; the degrees of these functions and of the differentials on the curves are related to numbers that appear in the Monstrous Moonshine. The idea for relating the curves is to factor out a (local) group action; this will be achieved by retaining only some of the functions in the affine rings of the curves, and manufacturing the relations they satisfy.

7.1. **Covering spaces of** $(3,7,8)$**, and** $(6,13,14,15,16)$ **curves.** We will relate to X_{12} certain covers and subcovers, with the goal of presenting observations on symmetries, acting on algebraic or transcendental functions, that appear to be related to the Monster.

First we should note that by sending y_7 and y_8 to y_{14} and y_{16}, the relations $f_{12,4}$, $f_{12,7}$ and $f_{12,9}$ for X_{12} map to f_{14}, f_{15}, and f_{16} for X_4, respectively. Hence X_{12} has a natural projection to X_4. More precisely, X_{12} is a double covering of X_4 as mentioned in the proof of Proposition 2.1.

The Jacobian $\mathcal{J}^{(4)}$ has 120 odd and 136 even θ characteristics for a total of $2^8 = 256$. Coble [C, Ch. 5] identified several group actions on the characteristics and Vakil [V] introduced an E_8 action. This sporadic group plays a role in the representation of the Monster [CN]. Through the double covering X_{12} over X_4, the basis of the tangent space of $\mathcal{J}^{(4)}$ is naturally embedded in those of $\mathcal{J}^{(12)}$, $\iota : T\mathcal{J}^{(4)} \hookrightarrow T\mathcal{J}^{(12)}$,

$$\frac{1}{2}\iota(\nu_1^{I;4}) = \nu_4^{I;12}, \quad \frac{1}{2}\iota(\nu_2^{I;4}) = \nu_6^{I;12}, \quad \frac{1}{2}\iota(\nu_3^{I;4}) = \nu_9^{I;12}, \quad \frac{1}{2}\iota(\nu_4^{I;4}) = \nu_{11}^{I;12}.$$

We have also have morphisms $X_4 \to X_6$ and $X_4 \to X_7$. Similarly X_{12} admits projections, from X_{12} to X_4, X_6, X_7 and the following two further curves X_2 and X_{30}:

(1) a hyperelliptic curve X_2 of genus two given by $f_{12,9} = y_{15}^2 - \hat{k}_2(x)k_3(x)$, and
(2) a $(6,13)$ singular curve X_{30} of genus $30 = g(\langle 6,13\rangle)$ given by $y_{13}^6 - \hat{k}_2(x)^3 k_2(x)^2 k_3(x)$.

As we showed in (4.6), we also have the natural projections from the affine curve $\hat{C}_{12}$ to the affine curve $C_{12} \subset X_{12}$, where $\hat{C}_{12} :=$ Spec $\hat{R}_{12}$ and $C_{12} :=$ Spec R_{12}. Since $\hat{R}_{12}$ is decomposed as $\mathbb{C}[x]$-algebra;

$$\hat{R}_{12} = \mathbb{C}[x, w_3]/(w_3^6 - k_3(x)) \otimes_{\mathbb{C}[x]} \mathbb{C}[x, w_2]/(w_2^6 - k_2(x)) \otimes_{\mathbb{C}[x]} \mathbb{C}[x, \hat{w}_2]/(\hat{w}_2^6 - \hat{k}_2(x)),$$

$\hat{C}_{12}$ is can be viewed as a fiber product of $C_1 \times_C C_0 \times_C \hat{C}_0$, where $C :=$ Spec $\mathbb{C}[x]$, $C_1 :=$ Spec $\mathbb{C}[x, w_3]/(w_3^6 - k_3(x))$, $C_0 :=$ Spec $\mathbb{C}[x, w_2]/(w_2^6 - k_2(x))$ and $\hat{C}_0 :=$ Spec $\mathbb{C}[x, \hat{w}_2]/(\hat{w}_2^6 - \hat{k}_2(x))$.

Additionally, $\hat{C}_{12}$ has a projection to $\hat{C}_4 :=$ Spec $\hat{R}_4$, $(\hat{C}_{12} \to \hat{C}_4)$ where

$$\hat{R}_4 := \mathbb{C}[x, \hat{y}_4]/(\hat{y}_4^3 - k_3(x)k_2(x))$$

by identifying $w_3 w_2$ with $\hat{y}_4$ as in (4.4), *i.e.*, $\hat{y}_4 := w_3 w_2$. This curve $\hat{C}_4$ is birational to the sextic affine curve $\hat{C}_4'$ for $w^3 = \prod_{i=0}^5 (z - c_i)$ which is transformed by,

$$z := \frac{1}{x - \tilde{c}_0} + c_0, \quad w := \frac{1}{\sqrt[3]{-\prod_{i=1}^5 (b_i - c_0)}} \frac{\hat{y}_4}{(z - c_0)^2}, \quad c_i := \frac{1}{b_i - \tilde{c}_0} + c_0,$$

giving local coordinates at ∞ for the Riemann surface $\hat{X}_4 = \hat{C}_4 \bigcup \hat{C}_4'$. The Jacobian $\hat{\mathcal{J}}^{(4)}$ of $\hat{X}_4$ also has 256 theta characteristics, the order of $(\mathbb{Z}/2\mathbb{Z})^8$, which Coble stated [C, p.210]. It is known that the reflection group of E_8 is related to algebraic curve of genus 4 and degree 6 via Del Pezzo surface and these theta characteristics [Man, Z]. The surface

$y^3 = x^5 + q^2$, which has the Du Val singularity and is also connected with E_8 and McKay correspondence [D, DPT, Sh, St].

7.2. **Numbers 10 and 19 in $L(6, 13, 14, 15, 16)$ and Norton Numbers.** Our first remark is that the complementary sequence of the sub-semigroup H_{12} of the non-negative integers $\mathbb{N}_0$ generated by $M_{12} := \{6, 13, 14, 15, 16\}$, namely the Weierstrass semigroup of X_{12} at ∞, is

$$L(H_{12}) := \{1, 2, 3, 4, 5, 7, 8, 9, 10, 11, 17, 23\},$$

while the Norton sequence is given by [FMN, McK, MS, No] (see Appendix A),

$$\{1, 2, 3, 4, 5, 7, 8, 9, 11, 17, 19, 23\}.$$

The sequences differ only by the elements 10 and 19. Using the covering $\hat{C}_{12}$ of C_{12}, we notice that the order of pole of the denominator of the holomorphic one form (4.7) of X_{12} decomposes as $10 + 19$:

$$\mathrm{wt}\left(\frac{1}{y_{13}y_{16}}\right) = \mathrm{wt}\left(\frac{1}{y_{14}y_{15}}\right) = \mathrm{wt}\left(\frac{1}{w_3^5 w_2^4 \hat{w}_2^3}\right) = 10 + 19,$$

because $\mathrm{wt}(w_3^2 w_2^2) = 10$ and $\mathrm{wt}(w_3^3 w_2^2 \hat{w}_2^3) = 19$ in $H^0(\hat{C}_{12}, \mathcal{O}_{\hat{C}_{12}})$. This points to the significance of the configurations of curves we consider.

7.3. **A similarity between $\sigma^{(12)}$ and replicable functions.** For a replicable function (see Appendix A)

$$f(q) = \frac{1}{q} + \sum_{i=1}^{\infty} c_i q^i,$$

the coefficients generated by a finite number of h_i's,

$$c_i \in \mathbb{Q}[h_1, h_2, h_3, h_4, h_5, h_7, h_8, h_9, h_{11}, h_{17}, h_{19}, h_{23}],$$

plays an essential role in the moonshine phenomena as in [FMN, McK, MS]. The Grunsky coefficients $h_{m,n}$ of $f(q)$ are defined by [Gr]

$$\sum_{m,n} h_{m,n} p^m q^n := \log\left(pq\frac{f(q) - f(p)}{p - q}\right),$$

and have the property,

$$h_{m,n} \in \mathbb{Q}[h_1, h_2, h_3, h_4, h_5, h_7, h_8, h_9, h_{11}, h_{17}, h_{19}, h_{23}],$$

Further it is known that the SL(2,$\mathbb{Z}$) action on replicable functions can be expressed in terms of Faber polynomials [MS]. Since τ-function solutions to the dKP hierarchy is also given by the Faber polynomial [CK], we hope that the sigma function $\sigma^{(12)}$ and its derivatives may provide information on replicable functions: an integrable system of KP-hierarchy can be typically solved in terms of theta, tau, sigma functions [EEG, Na2, Mul].

Remark 7.1.

(1) As mentioned in Remark 5.4, the $\sigma^{(12)}$ function over $\mathbb{C}^{12}$ is an entire function and is also expected to be expressed by the Schur function. If the moduli parameters $(b_1, \dots, b_7)$ are polynomials of p and q, it is expected that

$$\sigma^{(12)}(u) - S_{\mathcal{Y}_{12}}(T)|_{T_{(i)}=t_i} = \sum_{m,n} \hat{h}_{m,n} q^m p^n$$

where

$$\hat{h}_{m,n} \in \mathbb{Q}[\mathsf{t}_1, \mathsf{t}_2, \mathsf{t}_3, \mathsf{t}_4, \mathsf{t}_5, \mathsf{t}_7, \mathsf{t}_8, \mathsf{t}_9, \mathsf{t}_{10}, \mathsf{t}_{11}, \mathsf{t}_{17}, \mathsf{t}_{23}].$$

(2) $\sigma^{(12)}$ is a generalization of Weierstrass sigma function whereas the Weierstrass sigma function plays an important role in [HBJ]; if we consider a certain degenerate of the curve X_{12} associated with elliptic curves, it is expected that $\sigma^{(12)}$ might be written in terms of Weierstrass' elliptic σ.

(3) By the Riemann-Kempf theorem, using (4.9), a suitable derivative of σ, $\sigma_{\mathsf{t}_{i_1}, \cdots, \mathsf{t}_{i_k}}$, is a function of $\{\mathsf{t}_1, \mathsf{t}_2, \cdots, \mathsf{t}_{23-N^{(12)}(13-k)}\}$ over an open subset of the subvariety $\kappa^{-1}\mathcal{W}_k$ [MP12]. There the other t's of the hierarchy are functions of $\mathsf{t}_1, \mathsf{t}_2, \cdots, \mathsf{t}_{23-N^{(12)}(13-k)}$ [MP12]. This is analogous to the fact that for a suitable replicable function, every c_i belongs to a subring $\mathbb{Q}[h_1, \dots, h_\ell]$ of $\mathbb{Q}[h_1, h_2, h_3, h_4, h_5, h_7, h_8, h_9, h_{11}, h_{17}, h_{19}, h_{23}]$, and some of h's functions of the other h's, e.g., for j-function case, $c_i \in \mathbb{Q}[h_1, h_2, h_3, h_5]$ [Mah].

(4) For an (n, s) curve, algebraic solutions to the dKP equation exist [MP09]; it should be possible to generalize them to curves covered locally by affine patches such as X_4 and X_{12}. In other words, let us consider the case that $\hat{k}_2 = k_2$: Then $\phi_{11}^{(12)}\phi_9^{(12)} = (\phi_{10}^{(12)})^2$ and for $v := y_{14}/y_{15} = \dfrac{d}{d\mathsf{t}_2}\dfrac{\sigma_1^{(12)}}{\sigma_2^{(12)}} \Big/ \dfrac{d}{d\mathsf{t}_3}\dfrac{\sigma_1^{(12)}}{\sigma_2^{(12)}}$ over $\hat{u}(X_{12}) \subset \mathbb{C}^{12}$, we have a X_{12} curve solution of the dKP equation,

$$\frac{\partial}{\partial \mathsf{t}_3}\frac{\partial}{\partial \mathsf{t}_1} v + \frac{\partial}{\partial \mathsf{t}_1}\left(v \frac{\partial}{\partial \mathsf{t}_1} v \right) - 2 \frac{\partial}{\partial \mathsf{t}_2}\frac{\partial}{\partial \mathsf{t}_2} v = 0,$$

whose proof is the same as in [MP09]. It is expected that we might have the dKP hierarchy for $(\mathsf{t}_1, \mathsf{t}_2, \mathsf{t}_3, \mathsf{t}_4, \mathsf{t}_5, \mathsf{t}_7, \mathsf{t}_8, \mathsf{t}_9, \mathsf{t}_{10}, \mathsf{t}_{11}, \mathsf{t}_{17}, \mathsf{t}_{23})$. It is noted that J. McKay, and A. Sebbar considered the relation between the replicable functions and τ-functions of dKP hierarchy [MS].

A. APPENDIX: NORTON CONDITION

Let $\mathcal{A}$ be a filtered $\mathbb{Q}$ ring due to natural multiplication, $\mathcal{A} = \cup_i \mathcal{A}_i$ and $\mathcal{A}_j \mathcal{A}_i \subset \mathcal{A}_{i+j}$. Let us consider $q = \mathrm{e}^{2\pi\sqrt{-1}\tau}$ for $z \in H_+ := \{\tau \in \mathbb{C} \mid \Im\tau \geq 0\}$ and a function

$$f(q) = q^{-1} + h_1 q + h_2 q^2 + \cdots$$

where $h_j \in \mathcal{A}_j$. We also write it $f(\tau) = f(q)$.

The Grunsky coefficient $h_{m,n} \in \mathcal{A}_{m+n-1}$ of $f(q)$ is defined by [Gr]

$$\sum_{m,n} h_{m,n} p^m q^n := \log\left(pq\frac{f(q)-f(p)}{p-q}\right)$$

and Faber polynomial $F_{f,n}(f)$ [S, Fab] is defined by

$$F_{f,n}(f(q)) = \frac{1}{p^n} + n\sum_{m>0} h_{m,n} p^m,$$

where $h_{1,m} = h_m$. ¿From the definition of Grunsky coefficients, we have these property:

Lemma A.1.

$$h_{r,s} = h_{r+s-1} + \frac{1}{r+s}\sum_{m=1}^{r-1}\sum_{m=1}^{s-1}(n+m)h_{r+s-m-n-1}h_{m,n}.$$

These appear in the dispersionless KP hierarchy [CK].

The replicable functions are generalization of the the elliptic modular function $j(\tau)$, which is characterized by its expansion and the property under the action of Hecke operators T_n for every $n \geq 1$,

$$nT_n(j(\tau)) = \sum_{ad=n, 0\leq b<d} j\left(\frac{a\tau+b}{d}\right) = F_{f,n}(j(\tau)).$$

This gives $SL_2(\mathbb{Z})$ action on j.

The replicable functions were characterized by Norton, as having certain properties under the action of the Hecke operators, as members of the finite family of functions $\{f^{(a)}\}$ given by

$$\sum_{ad=n, 0\leq b<d} f^{(a)}\left(\frac{a\tau+b}{d}\right) = F_{f,n}(f(\tau)).$$

Norton considered these actions and characterized the properties of f or the coefficients $h_{m,1}$.

Definition A.2. *When for every case that $nm = rs$ and $(n,m) = (r,s)$, $h_{n,m} = h_{r,s}$, we say that $\mathcal{H}$ is replicable. The condition is called* Norton condition.

Example A.3. (1) $h_6 = h_{3,2} = h_4 + h_1 h_2$.
(2) $h_{12} = h_{3,4} = h_6 + h_1^2 h_2 + 2h_2 h_3 + h_1 h_4$
(3) $h_{10} = h_{5,2} = h_6 + h_1 h_4 + h_2 h_3$
(4) $h_{14} = h_{7,2} = h_8 + h_1 h_6 + h_2 h_5 + h_3 h_4$
(5) $h_{15} = h_{3,5} = h_7 + 2h_2 h_4 + h_3^2 + h_5 h_1 + h_1^2 h_3 + h_1 h_2^2$

The Norton condition is not consistent with the grading of $\mathcal{A}$ unless we modify the grading, e.g., $h_{-1} = 1$ with the weight -1.

Norton prove that [Cu],

Theorem A.4. *For a replicable function f with Fourier coefficients h_ℓ, every h_n belongs to $\mathbb{Q}[\mathcal{H}]$, where $\mathcal{H} := \{h_i \mid i \in \Phi\}$ and $\Phi := \{1, 2, 3, 4, 5, 7, 8, 9, 11, 17, 19, 23\}$.*

The set of the numbers $\ell = 1, 2, 3, 4, 5, 7, 8, 9, 11, 17, 19, 23$ is weakly symmetric except for the pair 4 and 10 or the pair 10 and 19 as shown in Table 4.

Table 4

0	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23
-	1	2	3	4	5	-	7	8	9	-	11	-	-	-	-	-	17	-	19	-	-	-	23
23	-	-	-	19	-	17	-	-	-	-	-	11	-	9	8	7	-	5	4	3	2	1	-
23	22	21	20	19	18	17	16	15	14	13	12	11	10	9	8	7	6	5	4	3	2	1	0

B. APPENDIX: COVERING OF CURVES

As the correspondence between an algebraic variety and a commutative ring, we have the well-known normalization theorem [Gri, p.5, p.68]:

Theorem B.1. *For any irreducible algebraic curve $X \subset P^2\mathbb{C}$, there exists a compact Riemann surface $\tilde{X}$ and a holomorphic mapping $s : \tilde{X} \to P^2\mathbb{C}$ such that $s(\tilde{X}) = X$ and s is injective on the inverse image of the set of smooth points of X. Further the Riemann surface is unique up to its isomorphism; if there are two Riemann surfaces $\tilde{X}$ and $\tilde{X}'$ given by normalizations X, there is a biholomorphic from $\tilde{X}$ to $\tilde{X}'$.*

Using this theorem, we have the following results:

B.1. Affine patches for the Riemann surface X_4.

As in the case of the (n, s) curves, X_4 can be covered with two affine patches (for the hyperelliptic case, cf. [Mum]).

Proposition B.2. *Without loss of generality, we assume that every b_a does not vanish. By letting $\underline{y}_7 := (y_7/x^3)$, $\underline{y}_8 := (y_8/x^3)$ and $\underline{x} := 1/x$,*

$$\begin{aligned} &\underline{f}_{14} := \underline{y}_7^2 - \underline{y}_8 \underline{x}\, \underline{k}_2(\underline{x}), \quad \underline{f}_{15} := \underline{y}_7 \underline{y}_8 - \underline{x}\, \underline{k}_2(\underline{x}) \underline{k}_3(\underline{x}), \\ &\underline{f}_{16} := \underline{y}_8^2 - \underline{y}_7 \underline{x}\, \underline{k}_3(\underline{x}), \end{aligned} \tag{B.1}$$

where $\underline{k}_3(x) = (1 - b_1 x)(1 - b_2 x)(1 - b_3 x)$ and $\underline{k}_2(x) = (1 - b_4 x)(1 - b_5 x)$,

$$\underline{\mathcal{U}}_4 := \begin{pmatrix} \frac{\partial \underline{f}_{14}}{\partial \underline{x}} & \frac{\partial \underline{f}_{14}}{\partial \underline{y}_7} & \frac{\partial \underline{f}_{14}}{\partial \underline{y}_8} \\ \frac{\partial \underline{f}_{15}}{\partial \underline{x}} & \frac{\partial \underline{f}_{15}}{\partial \underline{y}_7} & \frac{\partial \underline{f}_{15}}{\partial \underline{y}_8} \\ \frac{\partial \underline{f}_{16}}{\partial \underline{x}} & \frac{\partial \underline{f}_{16}}{\partial \underline{y}_7} & \frac{\partial \underline{f}_{16}}{\partial \underline{y}_8} \end{pmatrix},$$

whose rank is 2 around $(\underline{x}, \underline{y}_7, \underline{y}_8) = (0, 0, 0)$.

Proof. Since $\underline{\mathcal{U}}_4$ is

$$\underline{\mathcal{U}}_4 = \begin{pmatrix} \underline{y}_8 (\underline{x}\, \underline{k}_2(\underline{x}))' & 2\underline{y}_7 & -\underline{x}\, \underline{k}_2(\underline{x}) \\ (\underline{x}\, \underline{k}_2(\underline{x}) \underline{k}_3(\underline{x}))' & \underline{y}_8 & \underline{y}_7 \\ \underline{y}_7 \underline{k}_3'(\underline{x}) & -\underline{k}_3(\underline{x}) & 2\underline{y}_8 \end{pmatrix}.$$

it is obvious that its rank is 2 at $(\underline{x}, \underline{y}_7, \underline{y}_8) = (0,0,0)$. For the case that $(\underline{x}, \underline{y}_7, \underline{y}_8) \neq (0,0,0)$, since the structure is the same as that of $\mathcal{U}_4$, its rank is also 2. $\square$

We can define

$$\underline{R}_4 := \mathbb{C}[\underline{x}, \underline{y}_7, \underline{y}_8]/(\underline{f}_{14}, \underline{f}_{15}, \underline{f}_{16}),$$

and its Spec $\underline{R}_4$. We regard that the curve X_4 whose affine patches are Spec R_4 and Spec $\underline{R}_4$ is non-singular.

B.1.1. *Homogeneous ring of* X_4. By partially following the arguments on the space curve in Pinkham [Pi], in this subsection, we mention relations among Spec R_4, X_4, a homogeneous ring, its Proj, the numerical semigroup H_4, and the monomial curve related to B_4.

The coefficient ring is given by $\mathfrak{C}^{(4)} := \mathbb{C}[\hat{b}_1, \hat{b}_2, \hat{b}_3, \hat{b}_4]$ and its two projections are introduced by

$$\varphi_b^{(4)} : \mathfrak{C}^{(4)} \to \mathbb{C} = \mathfrak{C}^{(4)}/(\hat{b}_1 - b_1, \hat{b}_2 - b_2, \hat{b}_3 - b_3, \hat{b}_4 - b_4),$$

and

$$\varphi_0^{(4)} : \mathfrak{C}^{(4)} \to \mathbb{C} = \mathfrak{C}^{(4)}/(\hat{b}_1, \hat{b}_2, \hat{b}_3, \hat{b}_4).$$

We are now considering

$$R_4^{(b,z)} := \mathfrak{C}^{(4)}[x, y_7, y_8, z]/(f_{14}^{(b,z)}, f_{15}^{(b,z)}, f_{16}^{(b,z)}),$$

where for $k_3^{(b,z)} := (x - \hat{b}_1 z^3)(x - \hat{b}_2 z^3)(x - \hat{b}_3 z^3)$ and $k_2^{(b,z)} := (x - \hat{b}_4 z^3)(x - \hat{b}_5 z^3)$,

$$f_{14}^{(b,z)}(x, y_7, y_8, z) := y_7^2 - y_8 k_2^{(b,z)}, \quad f_{15}^{(b,z)}(x, y_7, y_8, z) := y_7 y_8 - k_2^{(b,z)}(x) k_3^{(b,z)}(x),$$

$$f_{16}^{(b,z)}(x, y_7, y_8, z) := y_8^2 - y_7 k_3^{(b,z)}(x).$$

Then we have $\varphi_b^{(4)}(f_a^{(b)}(x, y_7, y_8, 1)) = f_a(x, y_7, y_8)$ and $\varphi_0^{(4)}(f_a^{(b)}) = f_a^{(Z)}$ for $a = 14, 15, 16$. By letting $R_4^{(b)} := R_4^{(b,z)}/(z-1)$, we have natural homomorphisms,

$$\begin{tikzcd}
\mathrm{Spec}\ B_4 \arrow[r] \arrow[d] & \mathrm{Spec}\ R_4^{(b)} \arrow[d] & \mathrm{Spec}\ R_4 \arrow[l] \arrow[d] \\
\mathrm{Spec}\ \mathbb{C} \arrow[r] & \mathrm{Spec}\ \mathfrak{C}^{(4)} & \mathrm{Spec}\ \mathbb{C} \arrow[l]
\end{tikzcd}.$$

This shows that Spec R_4 and Spec B_4 are fibers over the moduli space of pointed curves with given Weierstrass semigroup H_4. $R_4^{(b)}$ is related to the moduli space $\mathcal{M}_4$ of $(3,7,8)$ curves [Pi].

We deal with the projective space by defining $R_4^{(z)} := \varphi_b^{(4)} R_4^{(b,z)}$. Here z^3 has the same weight as x. The weighted homogeneous ideal in $R_4^{(z)}$ due to $\mathbb{G}_m^{(4)}$ provides Proj $R_4^{(z)}$. We have

$$R_4^{(z)}/(z-1) \approx R_4, \quad R_4^{(z)}/(z) \approx B_4.$$

Thus we have the unique ∞ as a ramified point by letting $z = 0$, which recovers (3.1). More precisely, Z_3, Z_7 and Z_8 correspond to z^3/x, z^7/y_7, and z^8/y_8 whereas $\underline{x}$, $\underline{y}_7$ and

$\underline{y}_8$ correspond to z^3/x, y_7z^2/x^3, and y_8z/x^3; for example, the relation $f_{14}^{(b,z)}(x, y_7, y_8, z)$ becomes

$$\begin{aligned} f_{14}^{(b,z)}(x, y_7, y_8, z) &= \frac{y_7^2 y_8 x^2}{z^{14}} \left(\left(\frac{z^8}{y_8}\right) \left(\frac{z^3}{x}\right)^2 - \left(\frac{z^7}{y_7}\right)^2 \left(1 - \hat{b}_4 \frac{z^3}{x}\right) \left(1 - \hat{b}_5 \frac{z^3}{x}\right) \right) \\ &= \frac{x^6}{z^4} \left(\left(\frac{y_7 z^2}{x^3}\right)^2 - \left(\frac{y_8 z}{x^3}\right) \left(\frac{z^3}{x}\right) \left(1 - \hat{b}_4 \frac{z^3}{x}\right) \left(1 - \hat{b}_5 \frac{z^3}{x}\right) \right), \end{aligned}$$

around $z = 0$. Hence we could identify Proj $R_4^{(z)}$ with X_4 and the unique infinity point ∞ as $z = 0$ point. z can be regarded as element of $\mathbb{G}_m^{(4)}$.

The following relations hold, for functions w_i which are algebraic over the ring R_4, parametrizations of y_7 and y_8 by

$$y_7 = w_3 w_2^2, \quad y_8 = w_3^2 w_2, \tag{B.2}$$

where

$$w_2^3 = k_2, \quad w_3^3 = k_3. \tag{B.3}$$

Since $\tilde{R}_4 := \mathbb{C}[x, w_2, w_3]/(w_2^3 - k_2(x), w_3^3 - k_3(x)), = \mathbb{C}[x, w_2]/(w_2^3 - k_2(x)) \otimes_{\mathbb{C}[x]} \mathbb{C}[x, w_3]/(w_3^3 - k_3(x))$, its spectrum has a fiber structure.

B.2. **Riemann surface X_{12} and affine curves.** Let us consider another affine curve of X_{12} rather than one in Section 4. As for the $(3, 7, 8)$ curve, we can parametrize this curve with two smooth charts; we construct the one that contains ∞:

Proposition B.3. *Without loss of generality, we assume that every b_a does not vanish. Let us define $\underline{y}_a = y_a/x^3$ for $a = 13, 14, 15, 16$.*

$$\begin{aligned} &\underline{f}_{12,1} := \underline{y}_{13}^2 - \underline{x}\,\hat{\underline{k}}_2(\underline{x})\underline{y}_{14}, \quad \underline{f}_{12,2} := \underline{y}_{13}\underline{y}_{14} - \underline{x}\,\underline{k}_2(\underline{x})\underline{y}_{15}, \quad \underline{f}_{12,3} := \hat{\underline{k}}_2(x)\underline{y}_{14}^2 - \underline{y}_{13}\underline{y}_{15}\underline{k}_2(\underline{x}) \\ &\underline{f}_{12,4} := \underline{y}_{14}^2 - \underline{x}\,\underline{k}_2(\underline{x})\underline{y}_{16}, \quad \underline{f}_{12,5} := \underline{y}_{13}\underline{y}_{16} - \underline{y}_{14}\underline{y}_{15}, \quad \underline{f}_{12,6} := \underline{y}_{15}^2 - \underline{x}\,\hat{\underline{k}}_2(\underline{x})\underline{k}_3(\underline{x}), \\ &\underline{f}_{12,7} := \underline{y}_{14}\underline{y}_{16} - \underline{x}\,\underline{k}_2(\underline{x})\underline{k}_3(\underline{x}), \quad \underline{f}_{12,8} := \underline{y}_{15} y_{16} - \underline{k}_3(\underline{x})\underline{y}_{13}, \quad \underline{f}_{12,9} := \underline{y}_{16}^2 - \underline{k}_3(\underline{x})\underline{y}_{14}, \end{aligned}$$

where $\hat{\underline{k}}_2(x) = (1 - b_6x)(1 - b_7x)$. For every $(\underline{x}, \underline{y}_{13}, \underline{y}_{14}, \underline{y}_{15}, \underline{y}_{16})$ which is zero of every $(\underline{f}_{12,a})_{a=1,\cdots,9}$, we have

$$\operatorname{rank} \underline{\mathcal{U}}_{12} = 4, \quad \underline{\mathcal{U}}_{12} := \left(\left(\tfrac{\partial}{\partial \underline{x}} \underline{f}_{12,a} \quad \tfrac{\partial}{\partial \underline{y}_{13}} \underline{f}_{12,a} \quad \tfrac{\partial}{\partial \underline{y}_{14}} \underline{f}_{12,a} \quad \tfrac{\partial}{\partial \underline{y}_{15}} \underline{f}_{12,a} \quad \tfrac{\partial}{\partial \underline{y}_{16}} \underline{f}_{12,a} \right)_{a=1,\cdots,9} \right).$$

around $(\underline{x}, \underline{y}_{13}, \underline{y}_{14}, \underline{y}_{15}, \underline{y}_{16}) = (0, 0, 0, 0, 0)$.

Proof. $\underline{\mathcal{U}}_{12}$ is

$$\begin{pmatrix}
-(\underline{x}\,\underline{\hat{k}}_2)'\underline{y}_{14} & 2\underline{y}_{13} & -\underline{\hat{k}}_2 & & \\
-(\underline{x}\,\underline{k}_2)'\underline{y}_{15} & \underline{y}_{14} & \underline{y}_{13} & -\underline{x}\,\underline{k}_2 & \\
(\underline{x}\,\underline{\hat{k}}_2)'\underline{y}_{14}^2 - \underline{y}_{13}\underline{y}_{15}\underline{x}\,\underline{k}_2' & -\underline{y}_{15}\underline{x}\,\underline{k}_2 & 2\underline{x}\,\underline{\hat{k}}_2\underline{y}_{14} & -\underline{y}_{13}\underline{x}\,\underline{k}_2 & \\
-(\underline{x}\,\underline{k}_2)'\underline{y}_{16} & & 2\underline{y}_{14} & & \underline{k}_2 \\
 & \underline{y}_{16} & -\underline{y}_{15} & -\underline{y}_{14} & \underline{y}_{13} \\
-(\underline{x}\,\underline{\hat{k}}_2\underline{k}_3)' & & & 2\underline{y}_{15} & \\
-(\underline{x}\,\underline{k}_2\underline{k}_3)' & & \underline{y}_{16} & & \underline{y}_{14} \\
-\underline{k}_3'\underline{y}_{13} & -\underline{k}_3 & & \underline{y}_{16} & \underline{y}_{15} \\
-\underline{k}_3'\underline{y}_{14} & & -\underline{k}_3 & & 2\underline{y}_{16}
\end{pmatrix}.$$

At $(\underline{x}, \underline{y}_{13}, \underline{y}_{14}, \underline{y}_{15}, \underline{y}_{16}) = (0,0,0,0,0)$, it is obvious that its rank is 4. For the case that $(\underline{x}, \underline{y}_{13}, \underline{y}_{14}, \underline{y}_{15}, \underline{y}_{16}) \neq (0,0,0,0,0)$, since the structure is the same as that of $\mathcal{U}_{12}$, its rank is also 4. ☐

The scheme related to Spec R_{12} and Spec $\underline{R}_{12}$ is denoted by X_{12}. It has a unique ∞ point as $1/x = 0$.

B.2.1. *Homogeneous ring of* X_{12}. As mentioned in Section 4, we partially follow the arguments on the space curve by Pinkham [Pi] and describe X_{12} more precisely. The coefficient ring is given by

$$\mathfrak{C}^{(12)} := \mathbb{C}[\hat{b}_1, \hat{b}_2, \hat{b}_3, \hat{b}_4, \hat{b}_5, \hat{b}_6, \hat{b}_7]$$

and its two projections are introduced by

$$\varphi_b^{(12)} : \mathfrak{C}^{(12)} \to \mathbb{C} = \mathfrak{C}^{(12)}/(\hat{b}_1 - b_1, \hat{b}_2 - b_2, \hat{b}_3 - b_3, \hat{b}_4 - b_4, \hat{b}_5 - b_5, \hat{b}_6 - b_6, \hat{b}_7 - b_7),$$

and

$$\varphi_0^{(12)} : \mathfrak{C}^{(12)} \to \mathbb{C} = \mathfrak{C}^{(12)}/(\hat{b}_1, \hat{b}_2, \hat{b}_3, \hat{b}_4, \hat{b}_5, \hat{b}_6, \hat{b}_7).$$

We are now considering

$$R_{12}^{(b,\hat{z})} := \mathfrak{C}^{(12)}[x, y_{13}, y_{14}, y_{15}, y_{16}, \hat{z}]/\mathcal{P}^{(b,\hat{z})},$$

where for $k_3^{(b,\hat{z})} := (x - \hat{b}_1\hat{z}^6)(x - \hat{b}_2\hat{z}^6)(x - \hat{b}_3\hat{z}^6)$, $k_2^{(b,\hat{z})} := (x - \hat{b}_4\hat{z}^6)(x - \hat{b}_5\hat{z}^6)$, $\hat{k}_2^{(b,\hat{z})} := (x - \hat{b}_6\hat{z}^6)(x - \hat{b}_7\hat{z}^6)$,

$$\mathcal{P}^{(b,\hat{z})} := (f_{12,1}^{(b)}, f_{12,2}^{(b)}, f_{12,3}^{(b)}, f_{12,4}^{(b)}, f_{12,5}^{(b)}, f_{12,6}^{(b)}, f_{12,7}^{(b)}, f_{12,8}^{(b)}, f_{12,9}^{(b)}),$$

$$\begin{aligned}
f_{12,1}^{(b,\hat{z})} &:= y_{13}^2 - \hat{k}_2^{(b,\hat{z})} y_{14}, \quad f_{12,2}^{(b,\hat{z})} := y_{13}y_{14} - k_2^{(b,\hat{z})} y_{15}, \quad f_{12,3}^{(b,\hat{z})} := \hat{k}_2^{(b,\hat{z})}(x) y_{14}^2 - y_{13}y_{15}k_2^{(b,\hat{z})}(x) \\
f_{12,4}^{(b,\hat{z})} &:= y_{14}^2 - k_2^{(b,\hat{z})} y_{16}, \quad f_{12,5}^{(b,\hat{z})} := y_{13}y_{16} - y_{14}y_{15}, \quad f_{12,6}^{(b,\hat{z})} := y_{15}^2 - \hat{k}_2^{(b,\hat{z})}(x)\hat{k}_3^{(b,\hat{z})}(x), \\
f_{12,7}^{(b,\hat{z})} &:= y_{14}y_{16} - k_2^{(b,\hat{z})}(x)k_3^{(b,\hat{z})}(x), \quad f_{12,8}^{(b,\hat{z})} := y_{15}y_{16} - k_3^{(b,\hat{z})}(x)y_{13}, \\
f_{12,9}^{(b,\hat{z})} &:= y_{16}^2 - k_3^{(b,\hat{z})}(x)y_{14}.
\end{aligned}$$

Then we have $\varphi_b^{(12)}(f_{12,a}^{(b)}(x, y_{13}, y_{14}, y_{15}, y_{16}, 1)) = f_{12,a}$, and essentially $\varphi_0^{(12)}(f_{12,a}^{(b)}(x, y_{13}, y_{14}, y_{15}, y_{16}, 1)) = f_{12,a}^{(Z)}$ for $a = 1, 2, \cdots, 9$.

By letting $R_{12}^{(b)} := R_{12}^{(b,\hat{z})}/(\hat{z} - 1)$, we have natural homomorphisms,

$$\begin{tikzcd}
\text{Spec } B_{12} \arrow[r] \arrow[d] & \text{Spec } R_{12}^{(b)} \arrow[d] & \text{Spec } R_{12} \arrow[l] \arrow[d] \\
\text{Spec } \mathbb{C} \arrow[r] & \text{Spec } \mathfrak{C}^{(12)} & \text{Spec } \mathbb{C} \arrow[l]
\end{tikzcd} .$$

This shows that Spec R_{12} and Spec B_{12} are fibers over the moduli space corresponding to the numerical semigroup H_{12}. $R_{12}^{(b)}$ is related to the moduli space $\mathcal{M}_{12}$ of $(6, 13, 14, 15, 16)$ curves.

We should deal with the projective space by defining $R_{12}^{(\hat{z})} := \varphi_b^{(12)} R_{12}^{(b,\hat{z})}$. Here $\hat{z}^6$ has the same weight as x. Every homogeneous ideal in $R_{12}^{(\hat{z})}$ due to $\mathbb{G}_m^{(12)}$ provides Proj $R_{12}^{(\hat{z})}$, which is identified with X_{12}. We have

$$R_{12}^{(\hat{z})}/(\hat{z} - 1) \approx R_{12}, \quad R_{12}^{(\hat{z})}/(\hat{z}) \approx B_{12}.$$

Thus we have the unique ∞ as ramified point by letting $\hat{z} = 0$, which recovers (4.1), similarly to (3.1); Z_6, and Z_a $(a = 13, 14, 15, 16)$ correspond to $\hat{z}^6/x$ and $\hat{z}^a/y_a$ whereas $\underline{x}$, and $\underline{y}_a$ $(a = 13, 14, 15, 16)$ correspond to $\hat{z}^6/x$ and $\hat{z}^{18-a} y_a/x^3$.

References

[ACGH] E. Arbarello, M. Cornalba, P.A. Griffiths and J. Harris, Geometry of Algebraic Curves. Vol. I, Springer-Verlag, New York, 1985.

[A] T. Ayano, *Sigma Functions for Telescopic Curves*, arXiv1201.0644, .

[B1] H.F. Baker, Abelian functions. Abel's theorem and the allied theory of theta functions, Cambridge University Press, Cambridge, 1995, Reprint of the 1897 original.

[B2] H.F. Baker, *On the hyperelliptic sigma functions*, Amer. J. Math., **20** (1898) 301-384.

[B3] H.F. Baker, *On a system of differential equations leading to periodic functions*, Acta Math., **27** (1903) 135-156.

[BEL1] V.M. Buchstaber, V.Z. Enolskiĭ and D.V. Leĭkin, *Hyperelliptic Kleinian functions and applications*, in Solitons, geometry, and topology: on the crossroad, 1–33, Amer. Math. Soc. Transl. Ser. 2, 179, Amer. Math. Soc., Providence, RI, 1997.

[BEL2] V.M. Bukhshtaber, V.Z. Ènol'skiĭ and D.V. Leĭkin, *Rational analogues of Abelian functions*, Funct. Anal. Appl., **33** (1999) 83-94.

[BG1] Buchweitz, Ragnar-Olaf; Greuel, Gert-Martin Le nombre de Milnor equisingularite et deformations de singularites des courbes reduites. (French. English summary) C. R. Acad. Sci. Paris Seri. A et B 288 (1979), no. 1, A35-A38.

[BG2] Buchweitz, Ragnar-Olaf; Greuel, Gert-Martin The Milnor number and deformations of complex curve singularities. Invent. Math. 58 (1980), no. 3, 241-281.

[BLE1] V.M. Bukhshtaber, D.V. Leĭkin, and V.Z. Ènol'skiĭ, *σ-functions of (n, s)-curves*, Russian Math. Surveys, **54** (1999) 628-629.

[BLE2] V.M. Bukhstaber, D.V. Leĭkin, and V.Z. Ènol'skiĭ, *Uniformization of Jacobi manifolds of trigonal curves and nonlinear differential equations*, Funct. Anal. Appl., **34** (2000) 159-171.

[CK] R. Carroll and Y. Kodama, *Solutions of the dispersionless Hirota equation*, J. Phys. A: Math. & Gen., **28** (1995) 6373-6387.

[C] A. B. Coble, Algebraic geometry and theta functions, Amer. Math. Soc., New York, 1961, .

[CN] J. H. Conway and S. P. Norton, *Monstrous Moonshine*, Bull. London. Math. Soc., **11** (1979) 308-339.

[Cu] C. J. Cummins, *Some comments on replicable functions*, Modern trends in Lie algebra representation theory, (Queen's Univ., Kingston, ON, 1994) 48-55, Queen's paper in Pure and Appl. Math **94** (1994) .

[EEG] J.C. Eilbeck, V.Z. Enolskii and J. Gibbons, *Sigma, tau and Abelian functions of algebraic curves*, J. Phys. A: Math. Theor., **43** (2010) 455216 (20pp).

[EEL] J.C. Eilbeck, V.Z. Enolskii and D.V. Leykin, *On the Kleinian construction of abelian functions of canonical algebraic curves*, SIDE III—symmetries and integrability of difference equations (Sabaudia, 1998), 121–138, CRM Proc. Lecture Notes, 25, Amer. Math. Soc., Providence, RI, 2000.

[EEMOP1] J.C. Eilbeck, V.Z. Enolskii, S. Matsutani, Y. Ônishi, E. Previato, *Addition formulae over the Jacobian preimage of hyperelliptic Wirtinger varieties*, J. reine angew. Math., **619** (2008) 37–48.

[EEMOP2] J.C. Eilbeck, V.Z. Enolskii, S. Matsutani, Y. Ônishi, E. Previato, *Ableian Functions for Trigonal Curves of Genus Three*, Int. Math. Res. Not., **2007** (2007) 140,38.

[DPT] M. Demazure, H. Pinkham, and B. Teissier, *Séminaire sur les singularit'es des surfaces*, Lectre Notes in Math. 777, Springer-Verlag, Berlin-New York, 1980.

[D] A. Durfee, *Fifteen characterizations of rational double points and simple critical points*, L'Enseignement, **XXV** 1-2 (1979) 131-163.

[Fab] G. Faber, *Über polynomische Entwicklungen*, Math. Ann., **57** (1903) 389-408.

[FK] H. M. Farkas and I. Kra, Riemann Surfaces, second edition, Graduate Texts in Mathematics, Vol. 71. Springer-Verlag, Berlin-New York, 1991.

[Fay] J.D. Fay, Theta functions on Riemann Surfaces, Lecture Notes in Mathematics, Vol. 352. Springer-Verlag, Berlin-New York, 1973.

[FMN] D. J. Ford, J. McKay and S. P. Norton, *More on replicable functions*, Comm. in Alg., **22** (1994) 5175-5193.

[Her] J. Herzog, *Generators and relations of Abelian semigroup and semigroup ring*, Manu. Math., **3** (1970) 175-193.

[HBJ] F. Hirzebruch, T. Berger and R. Jung, Manifolds and Modular Forms, Vieweg, Bonn, 1992.

[Hi] N. Hitchin, *Harmonic spinors*, Advances in Math. 14 (1974), 1-55.

[G] D. Grand, *A generalization of a formula of Eisenstein*, Proc. Lond. Math. Soc., **62** (1991) 121-132.

[Gr] H.Grunsky, *Koeffizientenbedingungen für schlicht abbildende meromorphe Funktionen*, Math. Zeit., **45** (1939) 29-61.

[Gri] P. A. Griffiths, Introduction to Algebraic Curves, AMS Translations of Math. Monographs **76** 1989.

[K] F. Klein, *Über hyperelliptische Sigmafunctionen*, Math. Ann., **32** (1888) 351-380.

[K83] J. Komeda, *On Weierstrass points whose first non-gaps are four*, J. Reine. Angew. Math., **341** (1983), 68-86.

[K92] J. Komeda, *On Weierstrass points whose semigroups are generated by two elements*, Research Reports of Kanagawa Institute of Technology, **B-17** (1993), 211-217.

[K99] J. Komeda, *Existence of the primitive Weierstrass gap sequences on curves of genus 9*, Bol. Soc. Bras. Mat., **30** (1999), 125-137.

[K04] J. Komeda, *On 6-semigroups generated by 4-elements from which affine toric varieties can be constructed*, Research Reports of Kanagawa Institute of Technology **B-28** (2004), 79-85.

[KO04] J. Komeda and A. Ohbuchi *Weierstrass points with first non-gap four on a double covering of a hyperelliptic curve*, Serdica Math J., **30** (2004), 43-54; Corrigendum **32** (2006), 375-378.

[KO08] J. Komeda and A. Ohbuchi *Weierstrass points with first non-gap four on a double covering of a hyperelliptic curve II*, Serdica Math J., **34** (2008), 771-782.

[KO08a] J. Komeda and A. Ohbuchi *Existence of the non-primitive Weierstrass gap sequences on curves of genus 8*, Bull Braz Math. Soc. News Series, **39** (2008), 109-121.

[KS] D. Korotkin and V. Shramchenko, *On higher genus Weierstrass sigma-function*, Physica D, **241** (2012), 2086-2094.

[L] S. Lang, Introduction to algebraic and abelian functions, Second edition, Graduate Texts in Mathematics, 89. Springer-Verlag, New York-Berlin, 1982.

[Man] L. Manivel, *Configurations of lines and models of Lie algebras*, J. Alg., **304** (2006) 457-486.

[Mah] K. Mahler, *On a class of non-linear functional equations connected with modular functions*, J. Austral. Math. Soc., **22** (Ser. A) (1976) 65-120.

[Mat] H. Matsumura, Commutative ring theory, CSAM 8, , translated by M. Reid, Cambridge, Cambridge, 1986.

[Ma] S. Matsutani, *Hyperelliptic solutions of KdV and KP equations: re-evaluation of Baker's study on hyperelliptic sigma functions*, J. Phys. A, **34** (2001) 4721–4732.

[MK] S. Matsutani, *Sigma functions for a space curve (3, 4, 5) type with an appendix by J. Komeda*, arXiv:1112.4137 v2, (2012).

[MP08] S. Matsutani and E. Previato, Jacobi inversion on strata of the Jacobian of the C_{rs} curve $y^r = f(x)$, J. Math. Soc. Japan **60** (2008) 1009-1044.

[MP09] S. Matsutani and E. Previato, A class of solutions of the dispersionless KP equation, Phys. Lett., (2009) .

[MP12] S. Matsutani and E. Previato, Jacobi inversion on strata of the Jacobian of the C_{rs} curve $y^r = f(x)II$, to appear in J. Math. Soc. Japan.

[McK] J. McKay, *Essentials of Monstrous Moonshine*, Adv. Std. Pure Math., **32** (2001) 347-353.

[MS] J. McKay, and A. Sebbar, Fuchsian groups, automorphic forms and Schwarzians, Math. Ann. 318 (2000), no. 2, 255-275.

[Mul] M. Mulase, *Cohomological structure in soliton equations and Jacobian varieties*, J. Differential Geom. 19 (1984), no. 2, 403-430.

[Mum] D. Mumford, Tata Lectures on Theta II Jacobian theta functions and differential equations, Progress in Mathematics, 43. Birkhäuser Boston, Inc., Boston, MA, 1984.

[Na1] A. Nakayashiki, *On algebraic expansions of sigma functions for* (n, s) *curves*, Asian J. Math. **14** (2010), 175-212.

[Na2] A. Nakayashiki, *Sigma function as a tau function*, Int. Math. Res. Not. **2010** (2010) 373-94.

[No] S. P. Norton More on Moonshine, in Computational Group Theory eded M D. Atkinson, 1984 Academic Press, p.185-194.

[O] Y. Ônishi, *Determinant expressions for hyperelliptic functions*, Proc. Edinb. Math. Soc., **48** (2005) 1-38.

[Pi] H. C. Pinkham, *Deformation of algebraic varieties with* G_m *action*, Astérisque, **20** (1974) 1-131.

[Po] A. Polishchuk, Abelian Varieties, Theta Functions and the Fourier Transform, Cambridge, Cambridge, 2003.

[RG] J. C. Rosales and P. A. García-Sánchez, Numerical Semigroups, Springer-Verlag, New York-Berlin, 2008.

[R] G.F.B. Riemann, *Ueber das Verschwinden der ϑ-Functionen*, J. Reine Angew. Math., **65** (1866) 161-172.

[RV] Rim, Dock Sang; Vitulli, Marie A. Weierstrass points and monomial curves. J. Algebra 48 (1977), no. 2, 454-476.

[S] I. Schur, *On Faber polynomials*, Amer. J. Math., **67** (1945), 33-122.

[Sh] T. Shioda, *Construction of elliptic curves with high rank via the invariants of the Weyl groups*, J. Math. Soc. Japan, **43** (1991) 673-718.

[St] R. Stekolshchik, Notes on Coxeter Transfomrations and the McKay Correspondence, Springer-Verlag, New York-Berlin, 2008.

[V] R. Vakil, *Twelve points on the projective line, branched covers, and rational elliptic fibration*, Math. Ann., **320** (2001) 33-54.

[W] K. Weierstrass, *Zur Theorie der Abelschen Functionen*, J. Reine Angew. Math., **47** (1854) 289-306.

[Z] Y. G. Zarhin, *Del Pezzo surfaces of degree 1 and Jacobians*, Math. Ann., **340** (2008) 407-435.

Jiryo Komeda:
Department of Mathematics
Center for Basic Education and Integrated Learning,
Konagawa Institute of Technology,
Atsugi, 243-0292,
JAPAN.
e-mail: komeda@gen.kanagawa-it.ac.jp

Shigeki Matsutani:
8-21-1 Higashi-Linkan Minami-ku,
Sagamihara 252-0311,
JAPAN.
e-mail: rxb01142@nifty.com

Emma Previato:
Department of Mathematics and Statistics,
Boston University,
Boston, MA 02215-2411,
U.S.A.
e-mail: ep@bu.edu